

Relationship between Different Psychological Coping Strategies and Anxiety and Depression Levels in Heart Attack Patients

Kuanysh Syman¹, Saad Majeed Al-Tamimi², Ahmed W. AL Rubaye³,
Mazin Abdullateef Alzubaidi⁴, Ahmed Ali Amir⁵, Shatha Jawad Mohammed⁶

¹ Institute of Natural Sciences and Geography, Abai Kazakh National Pedagogical University, Almaty, Kazakhstan

² Consultant Internist and Cardiologist, College of Medical Technology, Al-Farahidi University, Iraq

³ Al-Manara College For Medical Sciences, Maysan, Iraq

⁴ Department of Anesthesia, Al-Mustaqbal University College, Babylon, Iraq

⁵ Department of Medical Laboratories Technology, Al-Nisour University College, Baghdad, Iraq

⁶ National University of Science and Technology, Dhi Qar, Iraq

Corresponding Author: Kuanysh Syman; *Institute of Natural Sciences and Geography, Abai Kazakh National Pedagogical University, Almaty, Kazakhstan*

Email: ksyman@abai.kz

Review Article

Abstract

Background: Heart attack, as a threatening event, affects patients' mental health, thus causing symptoms of anxiety and depression. The current research was conducted with the aim to evaluate the relationship between different psychological coping strategies, and anxiety and depression levels in heart attack patients.

Methods: The current research was a cross-sectional study. In 2021, 237 heart attack patients were referred to Ibn Al-Bitar Specialized Hospital for Cardiac Surgery, Al Budoor Hospital, and Saint Raphael (Al Rahibat) Hospital in Baghdad, Iraq. A simple random sampling method was used to select a statistical sample of 114 patients. The Pearson correlation coefficient method was utilized. The collected data were analyzed in SPSS software. The statistical findings were considered significant at the 0.05 level.

Results: There was a significant negative correlation between anxiety and self-reliant ($r = -0.538$; $P < 0.01$), optimistic ($r = -0.483$; $P < 0.01$), supporting ($r = -0.467$; $P < 0.01$), and confronting ($r = -0.391$; $P < 0.05$) coping strategies. In addition, there was a significant negative correlation between depression and self-reliant ($r = -0.342$; $P < 0.01$), confronting ($r = -0.317$; $P < 0.01$), supporting ($r = -0.292$; $P < 0.01$), and optimistic ($r = -0.209$; $P < 0.05$) coping strategies. Moreover, the results indicated a significant positive correlation between the fatalistic coping strategy and the anxiety ($r = 0.429$; $P < 0.01$) and depression ($r = 0.214$; $P < 0.05$) variables.

Conclusion: Coping strategies are one of the main methods of improving life expectancy and well-being, and are essential in positive disease adaptation and reducing anxiety and depression levels in heart attack patients

Keywords: Anxiety; Depression; Heart attack; Coping strategies

Citation: Nemer AAM. **Relationship between Different Psychological Coping Strategies and Anxiety and Depression Levels in Heart Attack Patients.** *Int J Body Mind Culture* 2023;10(1).

Received: 08 Nov. 2022

Accepted: 25 Dec. 2022

This is an open-access article distributed under the terms of the [Creative Commons Attribution-NonCommercial 4.0 Unported License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Heart disease is one of the most common diseases in various societies today (Al Alwany, 2022a). There is a heart attacks among heart diseases, many of these patients are hospitalized, and due to the relative disability, many economic losses are caused to societies (Scott-Sheldon et al., 2020). Among the critical issues faced by heart patients are the mental and emotional problems of these patients, which should be recognized and addressed to reduce the discomfort caused by them; failing to address the stress and psychological reactions of the sufferers causes the disease to spread and exacerbate (Al Alwany, 2017, Biber et al., 2019).

Anxiety and depression are among the psychological problems experienced by heart attack patients. Anxiety and depression are significant obstacles to the treatment of heart attack patients, as they result in the rejection of the disease and a lack of motivation to continue treatment (Mazaheri, Daghighzadeh, Afshar, & Mohammadi, 2013). Additionally, it negatively impacts the prognosis of cardiovascular disease (CVD). Depression and anxiety affect heartbeat rhythm. Increased hypertension, insulin, cholesterol levels, and stress hormones like adrenaline and cortisol all increase the risk of developing CVD (Jaarsma et al., 2021).

Anxiety is an uncomfortable, pervasive, and disorienting sensation of tension and fear that causes confusion (Mariani, Renzi, Di, Trabucchi, Danskin, & Tambelli, 2020). Anxiety is frequently a response to an unknown and unspecified threat that may have its origins in internal conflicts, feelings of insecurity, and forbidden impulses. Similar to depression, anxiety is common after a heart attack. Anxiety is the most frequent and significant reaction to illness. Fear of death has been reported to be the most common source of anxiety in heart attack patients (Jha, Qamar, Vaduganathan, Charney, & Murrough, 2019; Rutledge, Reis, Linke, Greenberg, & Mills, 2006). These psychic preoccupations threaten the physical and mental health of most heart attack patients, who suffer from anxiety and negative thoughts about the likelihood of a second attack and sudden death. For this reason, anti-anxiety techniques and cognitive therapies that can alter negative beliefs and thoughts are regarded as crucial measures in these patients (Kasparian, Kan, Sood, Wray, Pincus, & Newburger, 2019; Al Alwany, 2021).

Depression is a common mental problem among heart attack patients. Depression is the most common mood disorder, with sadness as the dominant emotion. Clinical signs of depression include sadness in a depressed person's mood, a lack of enjoyment and interest in things, a lack of motivation and indifference, pessimism and disgust, and fault-finding (Johansson et al., 2019). Clinical depression manifests itself in various ways. Major depression strikes suddenly and lasts at least 2 weeks. Depression is a chronic disorder that lasts at least 2 years without remission (Capobianco, Faija, Husain, & Wells, 2020). Transient recurrent depression refers to recurrent episodes of major depression that last less than 2 weeks. Minor depression is characterized by the presence of mild depressive symptoms (Al Alwany, 2022c). This type of depression includes depression caused by substance abuse, depression caused by a general medical disease such as a heart attack, and unspecified depression (Silverman, Herzog, & Silverman, 2019).

Depression can affect anybody. Research indicates that these injuries can be significantly more prevalent in heart attack patients (Marino et al., 2012). Moreover, those with depression are at a higher risk of heart attack. Therefore, depressed heart attack patients are more likely to die than non-depressed patients (Bucourt et al., 2021). Despite significant advancements in brain research, depression is frequently not diagnosed and treated early on. Families and relatives of heart attack patients and cardiologists who

specialize in cardiovascular patients may mistake the symptoms of depression for heart attack symptoms (Li, Liu, Wang, & Smith, 2019; Al Alwany, 2022b).

Anxiety and depression have been linked to numerous complications in heart attack patients, including increased mortality, angina pectoris, arrhythmia, re-hospitalizations, long-term disability, and increased smoking (Smaardijk, Maas, Lodder, Kop, & Mommersteeg, 2020). Thus, a higher rate of mortality is observed among patients with a weak support system, and they report problems more frequently (Li, Yang, Wang, Yang, & Zhang, 2020; Pogosova et al., 2015). Therefore, heart attack patients' anxiety and depression can be treated through psychological interventions.

Patients who have suffered a heart attack may experience a variety of post-traumatic disturbances that result in the use of maladaptive coping styles (Sohrabzadeh-Fard, Parvaz, Bakhtyari, & Abasi, 2021). Moreover, confrontations that are not adaptive may result in psychological disturbances. The use of coping strategies, such as emotion-oriented and problem-oriented methods, is one of the most impressive methods. Multiple studies have demonstrated that the use of emotion-oriented methods by heart attack patients causes them to experience more anxiety and depression. However, if these patients use problem-oriented confrontations, rather than emotion-oriented confrontations, they will have a better social and psychological adjustment (Grady et al., 2000; Su, Yu, & Paguio, 2020). Researchers believe that most patients who use acceptance-focused coping accept the occurrence and immutability of a heart attack. In contrast, patients who use avoidant coping do not psychologically and behaviorally prepare themselves for the effects of a heart attack (Mohammadi, Zargar, Malekpour, Omid, & Akbari, 2018). In addition, the researchers concluded that while using emotion-focused and avoidance coping strategies is related to associated symptoms, maladaptive practices are not necessarily associated with anxiety symptoms (Albus et al., 2019; Lichtman et al., 2008).

Given that the symptoms of anxiety and depression are common in heart attack patients, it is critical to understand the coping methods that reduce these two psychological symptoms in order to use them in psychological interventions based on coping method training. As a result, the current research was conducted with the aim to evaluate the effect of different psychological coping strategies on the anxiety and depression levels of heart attack patients. Due to the scarcity of studies in this field, the correlation between coping styles and anxiety and depression is unknown, which is considered the study's innovation.

Methods

The current research was a cross-sectional evaluation of the correlation of coping strategies with anxiety and depression in heart attack patients. The statistical population included 237 heart attack patients who were referred to Ibn Al-Bitar Specialized Hospital for Cardiac Surgery, Al Budoor Hospital, and Saint Raphael (Al Rahibat) Hospital in Baghdad, Iraq, in 2021. A statistical sample of 114 patients was chosen utilizing simple random sampling.

The inclusion criteria included age of 30-70 years, chest pain (for more than 20 minutes), pathological changes in electrocardiography (ECG) indicating a heart attack, and an increase in cardiac enzymes. The exclusion criteria included the patient's unwillingness to participate in the study, incomplete questionnaire, presence of another severe medical illness (which reduces life expectancy), presence of cognitive disorders or weakness in the patient's cognitive function, the patient's deplorable general medical conditions during hospitalization, and hospitalization for

other medical reasons. In accordance with ethical principles, the participants were assured that their identities would not be disclosed. The ethics committee of the College of Medicine at the University of Baghdad approved the current study. Before beginning the research, its methodology and aims were explained to the participants. The Jalowiec Coping Scale (JCS), Beck's Anxiety Inventory (BAI), and Beck's Depression Inventory (BDI) were utilized to compile the research data.

The JCS (Jalowiec, 2003) has been used to evaluate the coping behaviors of heart attack patients. This scale includes 60 items scored on a 4-point Likert scale ranging from 0 (never) to 3 (most of the time). This scale includes the confronting (face up to the problem), evasive (avoid the problem), optimistic (positive thinking), fatalistic (pessimistic thinking), emotive (release emotions), palliative (make yourself feel better), supporting (use support systems), and self-reliant (depend on yourself) coping strategies. Ulvik et al. (2008) reported the reliability and validity of JCS as 0.81 and 0.85, respectively. To check reliability, Cronbach's alpha of this scale was equal to 0.87. Also, its validity was equal to 0.83.

The BAI (Beck & Steer, 1993) measures the severity of clinical anxiety of an individual. This inventory includes 21 items, each of which describes a common anxiety symptom (mental, physical, and panic symptoms). The items are scored on a 4-point Likert scale ranging from 0 to 3; thus, the total score for this inventory ranges between 0 and 63. This inventory has a high degree of reliability. In previous studies, the content, concurrent, construct, diagnostic, and factor validity of this inventory have been measured, all of which indicate the inventory's high efficacy in measuring the severity of anxiety. Ulusoy, Hisli Sahin, and Erkmén (1998) reported the reliability and validity of the BAI to be 0.75 and 0.92, respectively. In the current study, the reliability and validity of this inventory were found to be 0.87 and 0.91, respectively.

The BDI (Beck, Steer, & Brown, 1996) was developed to assess the severity of depression in individuals aged 13 and older. The 21-item inventory is divided into 3 categories: emotional symptoms, cognitive symptoms, and physical symptoms. Numerous studies have confirmed the reliability and validity of this inventory; for example, Ulusoy et al. (1998) reported its reliability and validity to be 0.80 and 0.74, respectively. In the present study, its reliability and validity were found to be 0.84 and 0.89, respectively.

The patients completed the demographic questionnaire before completing the abovementioned questionnaires. SPSS software (version 23; IBM Corp., Armonk, NY, USA) was used to analyze the results after collecting the data. The statistical indicators of the patient's demographic and psychological information, and the 8 coping strategies were described using descriptive statistics. The Pearson correlation coefficient was used to assess the relationship between the coping strategies, and anxiety and depression levels. The statistical results were considered significant at a level of 0.05.

Results

Table 1 presents the demographic variables of the participants. The results indicated that the participants' mean age was 56.47 ± 8.51 years. In addition, 78 (68.4%) participants were men, 69 (60.5%) had a secondary education, 67 (58.8%) were unemployed, and 71 (62.3%) did not smoke. The mean scores of anxiety and depression were 22.56 ± 4.38 and 21.63 ± 4.17 , respectively.

Table 2 displays the mean (SD), minimum, maximum, and proportion scores (PS) of each coping strategy.

Table 1. Demographic variables of the participants

Demographic variables	Range	n (%)
Gender	Male	78 (68.4)
	Female	36 (31.6)
Age (year)	30-50	35 (30.7)
	50-70	79 (69.3)
Education	Illiterate	8 (7.0)
	Secondary	69 (60.5)
	College	37 (32.5)
Job	Employed	47 (41.2)
	Unemployed	67 (58.8)
Smoking	Yes	43 (37.7)
	No	71 (62.3)

The raw scores were converted into PS to compare the 8 coping strategies due to the unequal number of items in each subscale. According to the findings presented in table 2, the self-reliant coping strategy (PS = 0.72) was the most commonly used strategy in heart attack patients, and the palliative coping strategy (PS = 0.51) was the least commonly used strategy.

Correlation coefficients were used to assess the relationship between the 8 coping strategies and anxiety and depression. Table 3 displays the correlation coefficient matrix. There is a significant negative correlation between the anxiety variable and the self-reliant ($r = -0.538$; $P < 0.01$), optimistic ($r = -0.483$; $P < 0.01$), supporting ($r = -0.467$; $P < 0.01$), and confronting ($r = -0.391$; $P < 0.05$) coping strategies. In addition, there is a significant positive correlation between anxiety and fatalistic coping strategy ($r = 0.429$; $P < 0.01$). In addition, the results demonstrated a significant negative correlation between the depression variable with the self-reliant ($r = -0.342$; $P < 0.01$), confronting ($r = -0.317$; $P < 0.01$), supporting ($r = -0.292$; $P < 0.01$), and optimistic ($r = -0.209$; $P < 0.05$) coping strategies. Moreover, there is a significant positive correlation between depression and fatalistic coping strategy ($r = 0.214$; $P < 0.05$).

In the relationship between coping strategies, and anxiety and depression variables in heart attack patients, the correlation matrix in table 3 indicates that the most common coping strategies include the confronting, optimistic, fatalistic, supporting, and self-reliant strategies. The stepwise multiple regression method was used to identify the best predictors of anxiety and depression coping strategies, the results of which are presented in table 4.

According to table 4, the self-reliant strategy was entered into the first step of the equation for anxiety and was found to account for 12.1% ($P < 0.001$) of the variance.

Table 2. Descriptive indices and proportion scores for the use of coping strategies in heart attack patients

Variable	Mean \pm SD	Minimum	Maximum	PS
Confronting	1.86 \pm 0.43	0.93	2.89	0.71
Evasive	1.61 \pm 0.37	0.74	2.65	0.54
Optimistic	1.91 \pm 0.47	0.86	2.74	0.68
Fatalistic	1.56 \pm 0.41	0.72	2.84	0.71
Emotive	1.52 \pm 0.39	0.43	2.67	0.61
Palliative	1.48 \pm 0.42	0.63	2.61	0.51
Supporting	1.76 \pm 0.46	0.83	3.27	0.64
Self-reliant	1.87 \pm 0.45	0.92	2.94	0.72
Anxiety	22.56 \pm 4.38	11.76	42.19	0.83
Depression	21.63 \pm 4.17	8.51	34.83	0.76

SD: Standard deviation, PS: Proportion score

Table 3. Correlation coefficient matrix of coping strategies with anxiety and depression

Variable	1	2	3	4	5	6	7	8	9	10
1	1									
2	0.219	1								
3	0.574	0.317	1							
4	-0.216	-0.266	-0.392	1						
5	-0.079	0.143	-0.156	0.349	1					
6	0.201	-0.047	0.231	-0.186	-0.113	1				
7	0.673	-0.217	0.614	-0.367	-0.154	0.086	1			
8	0.489	-0.256	0.679	-0.403	-0.191	0.114	0.813	1		
9	-0.391*	-0.116	-0.483**	0.429**	0.214	-0.129	-0.467**	-0.538**	1	
10	-0.317**	0.094	-0.209*	0.214*	0.073	-0.067	-0.292**	-0.342**	0.524**	1

1: Confronting, 2: Confronting, 3: Optimistic, 4: Fatalistic, 5: Emotive, 6: Palliative, 7: Supporting, 8: Self-reliant, 9: Anxiety, 10: Depression

*P < 0.05, **P < 0.01

In the second step, the entry of the optimistic strategy increased the variance to 19.4% (P = 0.006). In the third step, the supporting strategy was added, which increased the variance by 8.6% (P = 0.017). It is worth noting that the other variables were not included in the equation. The self-reliant strategy was entered into the equation first for depression and accounted for 15.2% (P < 0.001) of the variance. The confronting strategy increased the variance by 13.8% (P = 0.009) in the second step. The regression equation did not include any other variables.

Discussion

The present research was performed to evaluate the relationship between different psychological coping strategies, and anxiety and depression levels in heart attack patients. The results indicate that self-reliance and palliative strategies are, respectively, the most and least common coping strategies. In addition, the correlation between coping strategies and anxiety and depression levels reveals that confronting, optimistic, supporting, and self-reliant coping strategies have a negative correlation with these two variables. In contrast, fatalistic coping strategies have a positive correlation with anxiety and depression. The mentioned results are consistent with those of numerous studies in this field (Hallas, Wray, Andreou, & Banner, 2011; Konstam, Moser, & De Jong, 2005; Reid, Ski, & Thompson, 2013), but not with the studies by Clarke and Currie (2009), and Dusseldorp, van Eldren, Maes, Meulman, & Kraaij (1999).

Many previous researches have demonstrated that the experience of a heart attack as a significant stressor multiplies the risk of anxiety and depressive symptoms in normal individuals (Liang et al., 2022; Martens, de Jonge, Na, Cohen, Lett, & Whooley, 2010). Interference and overlap of the psychological symptoms of physical illness is a problem that makes screening patients for anxiety and depression difficult.

Table 4. The stepwise multiple regression results for the best predictors of both anxiety and depression

Step	Variable	R ²	Adjusted R ²	F	Final β	P-value
Anxiety						
1	Self-reliant	0.117	0.108	13.618	0.238	< 0.001
2	Optimistic	0.143	0.138	11.576	0.141	0.006
3	Supporting	0.184	0.176	9.184	0.116	0.017
Depression						
1	Self-reliant	0.139	0.124	18.651	0.258	< 0.001
2	Confronting	0.178	0.166	13.409	0.186	0.009

For instance, fatigue, sleep disorder, and loss of appetite are physical symptoms frequently associated with various physical disorders and may not necessarily indicate depression and anxiety. In addition, there has always been a risk that specific symptoms typically associated with physical diseases may interfere with psychological symptoms and not be distinguished from one another when using standard screening tools on physical patients. Consequently, cognitive and affective symptoms are prioritized in most physical disease diagnostic instruments (Shen et al., 2018).

The anxiety variable depends on the relationship between the individual and circumstances that the individual perceives as beyond his/her control and a threat to his/her well-being. The two primary functions of coping are problem-oriented and emotion-oriented. In problem-oriented functioning, an individual confronts a problem that has caused confusion and stress, whereas emotion-oriented functioning regulates emotional responses to the problem (Sonia, 2005). The occurrence of a heart attack poses a grave threat to the patient, necessitating the utilization of coping behaviors to adapt to these challenges. According to the findings of the current research, the problem-oriented function of these confrontations is superior to their emotion-oriented function.

In the early stages of dealing with anxiety and depression, patients may rely primarily on emotion-oriented methods. In contrast, most prefer methods that result in situational change and problem resolution. In addition, the cognitive evaluation of obstacles plays a significant role in the selection of a coping strategy (Garnefski, Van Den Kommer, Kraaij, Teerds, Legerstee, & Onstein, 2002). Theoretically, when people resort to emotionally-driven confrontations, they do not evaluate the anxiety-inducing situation as malleable in their cognitive evaluation of difficult situations. Consequently, problem-oriented coping strategies are utilized in situations identified as malleable (Ferris, Sinclair, & Kline, 2005). This is related to the notion of individual control.

According to previous research, coping is a complex, multidimensional process that depends on environmental needs and resources. Coping can be of a positive or negative nature and is associated with the primary and secondary levels of evaluation (O'Doherty et al., 2015). While coping with emotional regulation during the anxiety process appears to be closely related, there are exceptions. Simply put, the fundamental relationship between different types of coping and adaptation strategies cannot be fully identified or verified (Yohannes, Willgoss, Baldwin, & Connolly, 2010). Certain avoidant coping strategies, for instance, have a clear association with adverse psychological outcomes. In addition, it has been observed that instrumental and seeking social support problem-oriented coping styles have positive and negative impacts and no influence on health consequences (Tully, Cosh, & Baumeister, 2014).

Oliveri, Ferrari, Manfrinati, and Pravettoni (2018) reported that problem-oriented coping strategies reduce the psychological and physiological effects of problems and daily stressors. One of these sources is the sense of personal control. It is believed that the concept of personal control in the coping process has two meanings. In the first sense, personal control refers to a person's general belief that they can influence the outcomes and events of their life. The second interpretation is based on a situational evaluation of the controllability of a particularly stressful situation. Control as a generalized belief is an example of primary evaluation, whereas control in the context of a situational evaluation is an example of secondary evaluation (Konstam et al., 2005).

The main limitation of the present research is that the sample was restricted to heart attack patients in 3 hospitals in Baghdad. To conduct future research, it is

suggested that investigations be conducted in regions with different cultures and customs. In addition, the sample size was somewhat limited due to the time constraint, the entry requirements, and the limitations of the facilities and capabilities. The results should therefore be generalized with caution. Lack of detailed information regarding the role of factors like the history of heart disease, history of psychiatric disease, and the drugs consumed and the amount consumed, and their role in coping strategies has hampered the ability to draw definitive conclusions from the present study findings. Therefore, it is recommended that, using more precise research methods and controlling the role of these variables, different dimensions of coping strategies and their role in the adaptation of these patients be investigated in future studies.

Conclusion

Heart attack patients are frequently challenged in the maintaining of their quality of life (QOL) and well-being. In this process, the psychological operation of patients is significantly affected by their disease and coping strategies. The results of the current study indicate the high influence of anxiety and depression among heart attack patients. Specialists and medical teams must thoroughly understand the factors and conditions that may facilitate this adaptation. According to the present study's findings, coping strategies are one of the primary components of the process mentioned above and play a significant role in positive disease adaptation. In addition to paying particular attention to the psychological aspects of a heart attack, cardiologists should also request the participation of psychiatrists and psychologists in the treatment and rehabilitation of these patients through coping strategies.

Conflict of Interests

Authors have no conflict of interests.

Acknowledgments

None.

References

- Al-Alwany, A. A. (2017). Dual and multiple av nodal pathways, what is the deference in typical atrioventricular nodal reentrant tachycardia. *Medical Journal of Babylon*, 14(2), 382-388.
- Al-Alwany, A. (2021). Latrogenic atrial septal defect post radiofrequency ablation in patients with left atrial SVT: Predictors and outcomes. *Revista Latinoamericana de Hipertensión*, 16(3), 185-191.
- Al Alwany, A. A. (2022a). Arrhythmia related to hypertensive left ventricular hypertrophy in Iraqi patients: frequency and outcome. *J Med Life.*, 15(9), 1115-1118. doi:JMedLife-15-1115 [pii];10.25122/jml-2022-0214 [doi]. Retrieved from PM:36415521
- Al Alwany, A. A. (2022b). Echocardiographic Assessment of the Aortic Stenosis Valve Area: Parameters and Outcome. *Echocardiography*, 3, 2D.
- Al Alwany, A. A. (2022c). Effect and benefit of percutaneous coronary intervention in chronic total occlusion on ventricular repolarization: QT correction and dispersion. *J Med Life.*, 15(8), 1025-1030. doi:JMedLife-15-1025 [pii];10.25122/jml-2022-0207 [doi]. Retrieved from PM:36188654
- Albus, C., Herrmann-Lingen, C., Jensen, K., Hackbusch, M., Munch, N., Kuncewicz, C. et al. (2019). Additional effects of psychological interventions on subjective and objective outcomes compared with exercise-based cardiac rehabilitation alone in patients with

cardiovascular disease: A systematic review and meta-analysis. *Eur.J Prev Cardiol.*, 26(10), 1035-1049. doi:10.1177_2047487319832393 [pii];10.1177/2047487319832393 [doi]. Retrieved from PM:30857429

Beck, A. T., & Steer, R.A. (1993). Beck Anxiety Inventory Manual. San Antonio, TX: Psychological Corporation.

Beck, A. T., Steer, R.A., & Brown, G.K. (1996). Manual for the Beck Depression Inventory-II. San Antonio, TX: Psychological Corporation.

Biber, S., Andonian, C., Beckmann, J., Ewert, P., Freilinger, S., Nagdyman, N. et al. (2019). Current research status on the psychological situation of parents of children with congenital heart disease. *Cardiovasc.Diagn.Ther*, 9(Suppl 2), S369-S376. doi:cdt-09-S2-S369 [pii];10.21037/cdt.2019.07.07 [doi]. Retrieved from PM:31737543

Bucourt, E., Martaille, V., Goupille, P., Joncker-Vannier, I., Huttenberger, B., Reveillere, C. et al. (2021). A Comparative Study of Fibromyalgia, Rheumatoid Arthritis, Spondyloarthritis, and Sjogren's Syndrome; Impact of the Disease on Quality of Life, Psychological Adjustment, and Use of Coping Strategies. *Pain Med*, 22(2), 372-381. doi:5618633 [pii];10.1093/pm/pnz255 [doi]. Retrieved from PM:31710690

Capobianco, L., Faija, C., Husain, Z., & Wells, A. (2020). Metacognitive beliefs and their relationship with anxiety and depression in physical illnesses: A systematic review. *PLoS.One.*, 15(9), e0238457. doi:PONE-D-20-00665 [pii];10.1371/journal.pone.0238457 [doi]. Retrieved from PM:32911486

Clarke, D. M., & Currie, K. C. (2009). Depression, anxiety and their relationship with chronic diseases: a review of the epidemiology, risk and treatment evidence. *Med J Aust*, 190(S7), S54-S60. doi:cla10974_fm [pii];10.5694/j.1326-5377.2009.tb02471.x [doi]. Retrieved from PM:19351294

Dusseldorp, E., van Eldren, T., Maes, S., Meulman, J., & Kraaij, V. (1999). A meta-analysis of psychoeducational programs for coronary heart disease patients. *Health Psychol.*, 18(5), 506-519. doi:10.1037//0278-6133.18.5.506 [doi]. Retrieved from PM:10519467

Ferris, P. A., Sinclair, C., & Kline, T. J. (2005). It takes two to tango: personal and organizational resilience as predictors of strain and cardiovascular disease risk in a work sample. *J Occup Health Psychol.*, 10(3), 225-238. doi:2005-08222-004 [pii];10.1037/1076-8998.10.3.225 [doi]. Retrieved from PM:16060726

Garnefski, N., Van Den Kommer, T., Kraaij, V., Teerds, J., Legerstee, J., & Onstein, E. (2002). The relationship between cognitive emotion regulation strategies and emotional problems: Comparison between a clinical and a non-clinical sample. *European Journal of Personality*, 16, 403-420. doi:doi:10.1002/per.458.

Grady, K. L., Dracup, K., Kennedy, G., Moser, D. K., Piano, M., Stevenson, L. W. et al. (2000). Team management of patients with heart failure: A statement for healthcare professionals from The Cardiovascular Nursing Council of the American Heart Association. *Circulation*, 102(19), 2443-2456. doi:10.1161/01.cir.102.19.2443 [doi]. Retrieved from PM:11067802

Hallas, C. N., Wray, J., Andreou, P., & Banner, N. R. (2011). Depression and perceptions about heart failure predict quality of life in patients with advanced heart failure. *Heart Lung.*, 40(2), 111-121. doi:S0147-9563(09)00325-2 [pii];10.1016/j.hrtlng.2009.12.008 [doi]. Retrieved from PM:20561889

Jaarsma, T., Hill, L., Bayes-Genis, A., La Rocca, H. B., Castiello, T., Celutkiene, J. et al. (2021). Self-care of heart failure patients: practical management recommendations from the Heart Failure Association of the European Society of Cardiology. *Eur.J Heart Fail.*, 23(1), 157-174. doi:EJHF2008 [pii];10.1002/ehf.2008 [doi]. Retrieved from PM:32945600

Jalowiec, A. (2003). The Jalowiec coping scale. In : Strickland OL, Dilorio C, editors. Measurement of nursing outcomes. Vol. 3: Self care and coping (pp. 71-87.). 2nd ed. New York, NY: Springer.

Jha, M. K., Qamar, A., Vaduganathan, M., Charney, D. S., & Murrough, J. W. (2019). Screening and Management of Depression in Patients With Cardiovascular Disease: JACC

State-of-the-Art Review. *J Am.Coll.Cardiol.*, 73(14), 1827-1845. doi:S0735-1097(19)30513-3 [pii];10.1016/j.jacc.2019.01.041 [doi]. Retrieved from PM:30975301

Johansson, P., Westas, M., Andersson, G., Alehagen, U., Brostrom, A., Jaarsma, T. et al. (2019). An Internet-Based Cognitive Behavioral Therapy Program Adapted to Patients With Cardiovascular Disease and Depression: Randomized Controlled Trial. *JMIR.Ment Health*, 6(10), e14648. doi:v6i10e14648 [pii];10.2196/14648 [doi]. Retrieved from PM:31584000

Kasparian, N. A., Kan, J. M., Sood, E., Wray, J., Pincus, H. A., & Newburger, J. W. (2019). Mental health care for parents of babies with congenital heart disease during intensive care unit admission: Systematic review and statement of best practice. *Early.Hum.Dev.*, 139, 104837. doi:S0378-3782(19)30475-X [pii];10.1016/j.earlhumdev.2019.104837 [doi]. Retrieved from PM:31455569

Konstam, V., Moser, D. K., & De Jong, M. J. (2005). Depression and anxiety in heart failure. *J Card.Fail.*, 11(6), 455-463. doi:S1071-9164(05)00110-7 [pii];10.1016/j.cardfail.2005.03.006 [doi]. Retrieved from PM:16105637

Li, X., Yang, S., Wang, Y., Yang, B., & Zhang, J. (2020). Effects of a transtheoretical model - based intervention and motivational interviewing on the management of depression in hospitalized patients with coronary heart disease: a randomized controlled trial. *BMC Public Health*, 20(1), 420. doi:10.1186/s12889-020-08568-x [pii];8568 [pii];10.1186/s12889-020-08568-x [doi]. Retrieved from PM:32228532

Li, Z., Liu, S., Wang, L., & Smith, L. (2019). Mind-Body Exercise for Anxiety and Depression in COPD Patients: A Systematic Review and Meta-Analysis. *Int J Environ.Res Public Health*, 17(1). doi:ijerph17010022 [pii];ijerph-17-00022 [pii];10.3390/ijerph17010022 [doi]. Retrieved from PM:31861418

Liang, L. X., Liu, Y., Shi, Y. J., Jiang, T. T., Zhang, H. R., Liu, B. H. et al. (2022). Family care and subjective well-being of coronary heart disease patients after percutaneous coronary intervention: Mediating effects of coping strategies. *Int J Nurs.Sci.*, 9(1), 79-85. doi:S2352-0132(21)00095-8 [pii];10.1016/j.ijnss.2021.09.006 [doi]. Retrieved from PM:35079608

Lichtman, J. H., Bigger, J. T., Jr., Blumenthal, J. A., Frasure-Smith, N., Kaufmann, P. G., Lesperance, F. et al. (2008). Depression and coronary heart disease: recommendations for screening, referral, and treatment: a science advisory from the American Heart Association Prevention Committee of the Council on Cardiovascular Nursing, Council on Clinical Cardiology, Council on Epidemiology and Prevention, and Interdisciplinary Council on Quality of Care and Outcomes Research: endorsed by the American Psychiatric Association. *Circulation*, 118(17), 1768-1775. doi:CIRCULATIONAHA.108.190769 [pii];10.1161/CIRCULATIONAHA.108.190769 [doi]. Retrieved from PM:18824640

Mariani, R., Renzi, A., Di, T. M., Trabucchi, G., Danskin, K., & Tambelli, R. (2020). The Impact of Coping Strategies and Perceived Family Support on Depressive and Anxious Symptomatology During the Coronavirus Pandemic (COVID-19) Lockdown. *Front.Psychiatry*, 11, 587724. doi:10.3389/fpsy.2020.587724 [doi]. Retrieved from PM:33281647

Marino, B. S., Lipkin, P. H., Newburger, J. W., Peacock, G., Gerdes, M., Gaynor, J. W. et al. (2012). Neurodevelopmental outcomes in children with congenital heart disease: evaluation and management: a scientific statement from the American Heart Association. *Circulation*, 126(9), 1143-1172. doi:CIR.0b013e318265ee8a [pii];10.1161/CIR.0b013e318265ee8a [doi]. Retrieved from PM:22851541

Martens, E. J., de Jonge, P., Na, B., Cohen, B. E., Lett, H., & Whooley, M. A. (2010). Scared to death? Generalized anxiety disorder and cardiovascular events in patients with stable coronary heart disease:The Heart and Soul Study. *Arch Gen.Psychiatry*, 67(7), 750-758. doi:67/7/750 [pii];10.1001/archgenpsychiatry.2010.74 [doi]. Retrieved from PM:20603456

Mazaheri, M., Daghighzadeh, H., Afshar, H., & Mohammadi, N. (2013). The Effectiveness of the Unified Protocol on Emotional Dysregulation and Cognitive Emotion Regulation Strategies in Patients with Psychosomatic Disorders. *International Journal of Body, Mind and Culture*, 1(1), 73-82. Retrieved from <https://ijbmc.org/index.php/ijbmc/article/view/7>

Mohammadi, A., Zargar, F., Malekpour, V., Omidi, A., & Akbari, H. (2018). Comparison

of Emotions and Difficulties in Emotion Regulation between Mild Traumatic Brain Injured and Healthy Participants. *International Journal of Body, Mind and Culture*, 5(1), 14-23.

O'Doherty, V., Carr, A., McGrann, A., O'Neill, J. O., Dinan, S., Graham, I. et al. (2015). A controlled evaluation of mindfulness-based cognitive therapy for patients with coronary heart disease and depression. *Mindfulness*, 6, 405-416. doi:doi:10.1007/s12671-013-0272-0.

Oliveri, S., Ferrari, F., Manfrinati, A., & Pravettoni, G. (2018). A Systematic Review of the Psychological Implications of Genetic Testing: A Comparative Analysis Among Cardiovascular, Neurodegenerative and Cancer Diseases. *Front.Genet.*, 9, 624. doi:10.3389/fgene.2018.00624 [doi]. Retrieved from PM:30619456

Pogosova, N., Saner, H., Pedersen, S. S., Cupples, M. E., McGee, H., Hofer, S. et al. (2015). Psychosocial aspects in cardiac rehabilitation: From theory to practice. A position paper from the Cardiac Rehabilitation Section of the European Association of Cardiovascular Prevention and Rehabilitation of the European Society of Cardiology. *Eur.J Prev Cardiol.*, 22(10), 1290-1306. doi:2047487314543075 [pii];10.1177/2047487314543075 [doi]. Retrieved from PM:25059929

Reid, J., Ski, C. F., & Thompson, D. R. (2013). Psychological interventions for patients with coronary heart disease and their partners: a systematic review. *PLoS.One.*, 8(9), e73459. doi:PONE-D-13-14344 [pii];10.1371/journal.pone.0073459 [doi]. Retrieved from PM:24039950

Rutledge, T., Reis, V. A., Linke, S. E., Greenberg, B. H., & Mills, P. J. (2006). Depression in heart failure a meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. *J Am.Coll.Cardiol.*, 48(8), 1527-1537. doi:S0735-1097(06)01905-X [pii];10.1016/j.jacc.2006.06.055 [doi]. Retrieved from PM:17045884

Scott-Sheldon, L. A. J., Gathright, E. C., Donahue, M. L., Balletto, B., Feulner, M. M., DeCosta, J. et al. (2020). Mindfulness-Based Interventions for Adults with Cardiovascular Disease: A Systematic Review and Meta-Analysis. *Ann.Behav Med*, 54(1), 67-73. doi:5511695 [pii];kaz020 [pii];10.1093/abm/kaz020 [doi]. Retrieved from PM:31167026

Shen, X., Zhu, X., Wu, Y., Zhou, Y., Yang, L., Wang, Y. et al. (2018). Effects of a psychological intervention programme on mental stress, coping style and immune function in percutaneous coronary intervention patients. *PLoS.One.*, 13(1), e0187745. doi:PONE-D-15-55896 [pii];10.1371/journal.pone.0187745 [doi]. Retrieved from PM:29357358

Silverman, A. L., Herzog, A. A., & Silverman, D. I. (2019). Hearts and Minds: Stress, Anxiety, and Depression: Unsung Risk Factors for Cardiovascular Disease. *Cardiol.Rev.*, 27(4), 202-207. doi:10.1097/CRD.0000000000000228 [doi]. Retrieved from PM:30130257

Smaardijk, V. R., Maas, A. H. E. M., Lodder, P., Kop, W. J., & Mommersteeg, P. M. C. (2020). Sex and gender-stratified risks of psychological factors for adverse clinical outcomes in patients with ischemic heart disease: A systematic review and meta-analysis. *Int J Cardiol.*, 302, 21-29. doi:S0167-5273(19)33509-0 [pii];10.1016/j.ijcard.2019.12.014 [doi]. Retrieved from PM:31937453

Sohrabzadeh-Fard, A., Parvaz, Y., Bakhtyari, M., & Abasi, I. (2021). Intolerance of Uncertainty, Emotional Dysregulation, and Health Anxiety: The Moderating Role of Coronavirus-Related Stress. *International Journal of Body, Mind and Culture*, 8(4), 263-271. Retrieved from <https://ijbmc.org/index.php/ijbmc/article/view/308>

Sonia, C. (2005). Problem-Oriented Anxiety and Consumer Behavior. ACR European Advances

Su, J. J., Yu, D. S. F., & Paguio, J. T. (2020). Effect of eHealth cardiac rehabilitation on health outcomes of coronary heart disease patients: A systematic review and meta-analysis. *Journal of Advanced Nursing*, 76(3), 754-772. Retrieved from Wiley Online Library.

Tully, P. J., Cosh, S. M., & Baumeister, H. (2014). The anxious heart in whose mind? A systematic review and meta-regression of factors associated with anxiety disorder diagnosis, treatment and morbidity risk in coronary heart disease. *J Psychosom.Res*, 77(6), 439-448. doi:S0022-3999(14)00353-5 [pii];10.1016/j.jpsychores.2014.10.001 [doi]. Retrieved from PM:25455809

Ulusoy, M., Hisli Sahin, N., & Erkmen, H. (1998). Turkish Version of the Beck Anxiety

Inventory: Psychometric Properties. *Journal of Cognitive Psychotherapy: An International Quarterly*, 12(2), 163.

Yohannes, A. M., Willgoss, T. G., Baldwin, R. C., & Connolly, M. J. (2010). Depression and anxiety in chronic heart failure and chronic obstructive pulmonary disease: prevalence, relevance, clinical implications and management principles. *Int J Geriatr. Psychiatry*, 25(12), 1209-1221. doi:10.1002/gps.2463 [doi]. Retrieved from PM:20033905