


Article type:
Original Research

1 Department of Psychology, Qom Branch, Islamic Azad University, Qom, Iran.

2 Department of Education Science, Mashhad Branch, Islamic Azad University, Mashhad, Iran.

Corresponding author email address:
p.sabzevari@iaush.ac.ir



Article history:

Received 18 Sep 2025

Revised 24 Nov 2025

Accepted 12 Dec 2025

Published online 01 Jan 2026

How to cite this article:

Omidvar, F., & Sabzevari, P. (2026). Effect of Acceptance and Commitment Therapy on Anxiety and Alexithymia of Mothers Having Children with Insecure Attachment. International Journal of Body, Mind and Culture, 13(1), 71-79.



© 2025 the authors. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

Effect of Acceptance and Commitment Therapy on Anxiety and Alexithymia of Mothers Having Children with Insecure Attachment

Fatemeh. Omidvar¹, Parisa. Sabzevari^{2*}

ABSTRACT

Objective: Psychologically healthy mothers who create a happy and peaceful environment can raise children with a secure attachment style. This research investigates the impact of acceptance and commitment therapy on anxiety and alexithymia in mothers with children exhibiting insecure attachment styles.

Methods and Materials: The research employs a semi-experimental design with pre-test/post-test measures and control and experimental groups. The statistical population comprises mothers with children exhibiting insecure attachment styles in health, educational, therapeutic, and preschool centers in Tehran during the spring and summer of 2023. The data collection tools included Kappenberg and Halpern's attachment questionnaire (2006), Beck's anxiety scale questionnaire (1998), and the Toronto alexithymia scale questionnaire (1994). An intervention program utilizing Hadi Toroghi and Amiri Hashemi's (2021) acceptance and commitment therapy package was administered to the experimental group. This research employed the analysis of covariance method to examine the pre-test and post-test data with SPSS Software by Version 23.

Findings: The results indicated that the level of anxiety and alexithymia underwent significant changes following the intervention of ACT. Acceptance and commitment therapy was effective in reducing anxiety ($F=24.720$, $P=0.001$) and alexithymia ($F=0.256$, $P=0.0358$) in mothers with children exhibiting insecure attachment styles.

Conclusion: Counseling and psychotherapy centers should consider implementing these methods, particularly acceptance and commitment therapy, more widely and with a greater focus.

Keywords: Acceptance and Commitment Therapy, Alexithymia, Anxiety, Insecure Attachment.

Introduction

Children with a secure attachment style tend to develop long-term relationships built on trust and emotional connection as adults. Moreover, these individuals typically exhibit high levels of self-esteem, enjoy intimate relationships, seek social support, and feel comfortable sharing their feelings with others, children who find secure attachment feel comforted, and their needs are met by their parents and caregivers. People with secure attachment may not enter into relationships with individuals who have anxious or avoidant attachments due to being self-aware that their needs are not being fulfilled within the relationship (Feyzabadi et al., 2025; Zhao et al., 2023). Research has demonstrated that individuals with a secure attachment style often display healthy relationships with their mothers, intimacy, trust, affection for others, and positive social interactions (Moghadam, 2024) (Monfaredi et al., 2022). Attachment disorders are measured along two dimensions: avoidance (such as mistrust in others' intentions and excessive self-reliance) and anxiety (such as worrying about the unavailability of others when needed and anxiously searching for support and love) (Akbari & Elmi, 2017; Mikulincer & Shaver, 2012). Insecure attachment styles are associated with emotional dysregulation and trait hyperarousal (Çiçek Gümüş & Öncel, 2023). Individuals with an anxious insecure attachment style may exhibit physical and cognitive hyperarousal, anxiety in Mothers have harmed the mother-baby relationship and the mother's ability it reduces maternal role while those with an avoidant attachment style may display parasympathetic nervous system activity, anxious attachment, or insecure attachment, refers to a behavior pattern in relationships in which a person is constantly afraid of abandonment. In simpler terms, a person with anxious attachment is always worried that their emotional partner will leave them and therefore tries to control them. This type of attachment exists in both children and adults (Darvishinia, 2023; Marino et al., 2021).

Another variable that has been studied in individuals with generalized anxiety is alexithymia, which refers to the absence of words for feelings, people with alexithymia magnify normal physical stimuli, misinterpret physical signs of emotional arousal, exhibit

emotional helplessness through physical complaints, and seek to address physical symptoms in therapy. Emotional dyslexia is a complex phenomenon that includes difficulties in identifying and distinguishing between emotions and physical arousal related to emotions. It also involves challenges in expressing feelings to others, limited visualization abilities, and a tendency towards a more objective, pragmatic, and realistic thinking style (Askari & Karami, 2024; Wijk et al., 2023). If the negative emotions drain and the person cannot express his negative feelings to fully express, the psychological component of emotional expression systems and increased psychological distress, including anxiety and depression finds (Abdi Zarrin & Nikkhah Siruei, 2021). Research has shown that attachment style is associated with alexithymia, with the characteristics of alexithymia being more commonly observed in individuals with insecure attachment styles (Jiang et al., 2024; Zhao et al., 2023). Kraemer and Loader suggest that insecure attachment may hinder the learning of emotions and contribute to the development of alexithymia. Loas et al. (2015) demonstrated that insecure attachment relationships predict difficulties in identifying and expressing emotions. Consequently, a mother who experiences anxiety and issues such as alexithymia may struggle to maintain a healthy relationship with her child, potentially leading to an insecure attachment style.

Individuals with alexithymia face difficulties in recognizing, expressing, processing, and regulating emotions. Alexithymia is considered a deficit in emotional self-regulation (Lumley et al., 2005) and is a multifaceted construct characterized by difficulty in identifying feelings, differentiating between emotions, experiencing bodily arousal in response to emotional stimuli, describing emotions to others, limited power of visualization, and a practical and realistic cognitive style or objective thinking (Byrne et al., 2021).

Several studies suggest that the insecure relationship between mother and child, which ultimately leads to insecure attachment, may also serve as a foundation for anxiety in adolescence and adulthood. Wiltgen et al. (2015) reported that avoidant and anxious attachment styles are related to anxiety. Research on attachment in adulthood has revealed a negative relationship between secure attachment and indicators of psychological distress, such as anxiety (Gould et al., 2018). Hahs et al.

(2019) found that insecure attachment is associated with anxiety and internalizing problems in childhood. Anxiety is often viewed as a response to future threats.

Over the years, various psychological treatments have been developed to address psychological problems such as anxiety and its related factors like alexithymia. Currently, we are witnessing the emergence of the third generation of these treatment approaches, often referred to as acceptance-based models. Examples of these models include mindfulness-based cognitive therapy, dialectical behavior therapy, metacognitive therapy, acceptance and commitment therapy, and even cancer therapy (Eskandari et al., 2024). These treatments place emphasis on the cognitive process rather than changing the cognitive content, with the aim of enhancing the individual's psychological connection with their thoughts and feelings.

Acceptance and commitment therapy (ACT) is considered an emerging third-generation treatment approach that is particularly suitable for Iranian patients due to its incorporation of Eastern techniques. ACT is based on a philosophical theory known as "functional contextualism" and is grounded in the "theory of mental interfaces framework," which explores language and cognition. ACT focuses on six central processes that promote psychological flexibility: acceptance, defusion, self-as-context, present moment awareness, values, and committed action (Hayes et al., 2006). Cognitive flexibility is associated with reduced psychological distress in chronic diseases disabling (Karimian et al., 2023).

Psychologically healthy mothers who create a peaceful environment can foster a secure attachment style in their children. On the other hand, mothers who experience anxiety and issues such as alexithymia may contribute to the development of an insecure attachment style in their children. This research aims to identify effective ways to address maternal anxiety and alexithymia, with the ultimate goal of promoting healthy relationships between mothers and their children. Therefore, the study seeks to determine whether acceptance and commitment therapy can effectively alleviate anxiety and alexithymia in mothers who have children with an insecure attachment style.

To date, there have been no direct or indirect studies examining the impact of acceptance and commitment therapy on anxiety and alexithymia in mothers who have

children with an insecure attachment style. Therefore, this research aims to address this research gap and contribute a new insight into the effectiveness of acceptance and commitment therapy for reducing anxiety and alexithymia in mothers with children exhibiting insecure attachment.

Methods and Materials

The current research was a semi-experimental study employing a pre-test-post-test design with control and experimental groups. The study's statistical population consists of mothers with children exhibiting insecure attachment who attend health, educational, therapeutic, and preschool centers in Tehran during the spring and summer of 2023. Participants were selected through a screening process using the children's attachment questionnaire during the mid-period. Specifically, mothers who have children with high scores indicating insecure attachment were eligible for the study if they reported experiencing anxiety and alexithymia.

The inclusion criteria for participation in the research require a minimum education level of secondary education. Mothers who have children with insecure attachment and attend health, educational, therapeutic centers, and preschools are eligible to participate in the intervention sessions. The exclusion criteria for discontinuing participation in the research include mothers who have a chronic medical or psychiatric condition, use specific medications, fail to complete the research questionnaires, are absent for more than two sessions of the designated intervention, participate in similar treatment sessions concurrently or within the past six months, and lack willingness and consent to participate.

Sample Size

Using Cohen's table, the minimum sample size for each group, with a permissible error rate of 0.5, is 14 individuals. However, to ensure a sufficient return of the questionnaires, we have rounded this number up to 15 individuals for each group. As a result, the total number of observations is 30. Using random sampling with replacement, the study recruited 30 participants who were assigned to either the control (n=15) or experimental (n=15) group.

Instruments

Halpern and Kappenberg's attachment

questionnaire: The questionnaire designed by (Malik et al., 2015) measures the attachment of children in the middle age period of childhood, specifically between pre-primary and primary school ages (3 to 12 years). The questionnaire is completed by the mother and consists of 20 items. The initial reliability of the questionnaire has been reported as 0.85 based on internal correlation and 0.83 based on the split-half method in the study conducted by the creators of the questionnaire. In a test-retest repeatability study involving 23 children, the reliability of the questionnaire was found to be 0.79. The creators of the scale have reported Cronbach's alpha coefficients of 0.69, 0.63, 0.65, and 0.56 for positive adaptive developmental components, negative behaviors, emotional reactions, and avoidance of support from the attachment figure/caring person respectively. The construct validity of the questionnaire was examined using the Randolph Attachment Disorder Questionnaire (RADQ) with favorable results. (Kappenberg & Halpern, 2006) also demonstrated the favorable psychometric properties of this questionnaire. In the present research, the reliability of the attachment scale was found to be 0.695.

Beck Anxiety Inventory (BAI): The Beck Anxiety Inventory, created by Beck et al., (1988), is a self-report questionnaire specifically designed to measure the severity of clinical anxiety symptoms in individuals. This inventory was developed to assess anxiety intensity in both adolescents and adults. It consists of 21 statements, with each statement reflecting a symptom of anxiety. Responses are graded on a 4-point Likert scale ranging from 0 to 3, where 0 represents 'never,' 1 represents 'mild,' 2 represents 'moderate,' and 3 represents 'severe.' The internal consistency coefficient of the Beck Anxiety Inventory is reported to be 0.92, indicating good reliability. Test-retest reliability with a one-week interval is reported to be 0.75, and the correlation among its items ranges from 0.30 to 0.76. In this particular research, the inventory obtained a Cronbach's alpha coefficient of 0.73. The Beck Anxiety Inventory will be administered as both a pre-test and a post-test in this study. The reliability of the anxiety scale used in this research was found to be 0.716."

Toronto Alexithymia Scale Questionnaire: "The Toronto Alexithymia Scale Bagby et al., (1994) is a 20-question test consisting of three subscales: difficulty in identifying feelings, difficulty in describing feelings, and

objective thinking. Responses are rated on a 5-point Likert scale ranging from 1 ('I completely disagree') to 5 ('I completely agree'). The psychometric properties of the Toronto-20 alexithymia scale have been examined and confirmed in several studies (Bagby et al., 1994; Marsero et al., 2011). The Farsi version of the Toronto-20 Alexithymia Scale demonstrated good internal consistency, with Cronbach's alpha coefficients of 0.85 for total alexithymia, 0.82 for difficulty in identifying feelings, 0.75 for difficulty in describing feelings, and 0.72 for objective thinking. Retest reliability of the Toronto-20 alexithymia scale was confirmed in a sample of 67 individuals over two occasions with an interval of four weeks, with coefficients ranging from 0.80 to 0.87 for total alexithymia and different subscales (Rostamifar & Sajjadi). In this research, the Toronto Alexithymia test was administered as both a pre-test and post-test."

The independent variable, interventions of acceptance and commitment therapy, were implemented for the experimental group over a period of 9 sessions, with each session lasting 2 hours and conducted once per week. The research utilized seven therapeutic protocols from the book 'Acceptance and Commitment Therapy for Anxiety Disorders' by George Eifert and John Forsyth, as well as the 'Handbook of Acceptance and Commitment Therapy for Depression and Anxiety' by (MEHRABIZADEH et al., 2009).

Analysis

This research employed the analysis of covariance method to examine the pre-test and post-test data. The statistical analysis method used for the anxiety variable was analysis of variance, as there were no sub-components. For the Alexithymia variable, the statistical analysis method used was analysis of covariance, due to the presence of sub-components.

Ethics

This research obtained approval from Islamic Azad University for adhering to IR.IAU.QOM.REC.1402.125. Written consent was obtained from the participants to participate in the research.

Findings and Results

The 38% of respondents are under 25 years old, 40% are between 25-35 years old, 17% are between 35-45 years old, and the remaining 5% are over 45 years old.

According to the results presented in Table 1,

However, the mean scores (and standard deviations) of these variables in the control group did not show a noticeable change from the pre-test stage to the post-test stage.

Assumption is confirmed for all dependent variables in this study. Hence, the analysis of variance method can

be used to test the hypotheses due to the random assignment of participants to the experimental and control groups and the adequate sample size. Moreover, as the F-test is resistant to moderately heterogeneous variances, particularly when sample sizes are equal, it is appropriate to use the analysis of variance test.

Table 1

Descriptive data of dimensions of alexithymia and anxiety by group and measurement stage

Dependent variable	Group	Number	Pre-test	Post-test
			M \pm SD	M \pm SD
Anxiety	Intervention	15	34.3 \pm 11.60	72.1 \pm 65.42
	Control	15	36.64 \pm 25.60	77.3 \pm 13.60
Difficulty in identifying feelings,	Experiment	15	57.2 \pm 14.23	66.2 \pm 54.17
	Control	15	98.1 \pm 86.22	88.1 \pm 40.22
Difficulty in describing feelings	Experiment	15	66.2 \pm 75.19	17.2 \pm 19.16
	Control	15	22.2 \pm 18.19	45.2 \pm 38.19
Objective thinking	Experiment	15	79.2 \pm 68.22	56.2 \pm 72.17

The acceptance and commitment intervention had a significant effect on at least one variable of anxiety and alexithymia in the post-test stage ($P<0.01$, $F=513.702$). Specifically, the acceptance and commitment therapy was found to improve the anxiety and alexithymia of

mothers who participated in this study. Therefore, it can be concluded that the intervention based on acceptance and commitment was effective in reducing mothers' anxiety and alexithymia.

Table 2

Results of the analysis of variance of the measurement of anxiety in two stages of implementation

Research variables	Change sources	F	df	P-value	Effect size
Anxiety	Group	720.24	1	001.0	371.0
	Time	611.5	28	007.0	211.0
	Group & time	380.6		001.0	371.0

According to the results presented in Table 2, the intervention method (acceptance and commitment therapy) had a significant effect on anxiety scores in the three stages of measurement ($F=24.720$; $Sig=0.001$) for both the experimental and control groups. This indicates

that the acceptance and commitment therapy significantly improved participants' anxiety levels. Table 3 provides a two-by-two comparison of the adjusted means for the different stages of the anxiety test (pre-test, post-test).

Table 3

Main results of the covariance test of the effect of acceptance and commitment therapy on mothers' alexithymia

Source	df	Mean Square	F	Sig	Partial Eta Squared
Pre-test	1	18.60	0.895	0.036	0.029
Group	1	4.08	0.256	0.0358	0.016
Error	27	19.20			

Table 3 presents the major results of the covariance test, showing that a value of sig lower than 0.05 indicates the significance of the test. As shown in the group

variable row, the number is significant and equals 0.0358. This indicates a significant difference between the control and experimental groups in alexithymia. The

effect size is represented by the eta coefficient, which is visible in the Partial Eta Squared column. Multiplying this number by 100 gives us the percentage of variance in the dependent variable (alexithymia) explained by the same grouping variable. In this case, multiplying the number 0.016 by 100 yields 1.6, indicating that the independent variable explains 1.6% of the variance in the dependent variable (alexithymia).

Discussion and Conclusion

The aim of the study in this research was to investigate the effect of acceptance and commitment therapy on anxiety and alexithymia in mothers of children with insecure attachment. The results indicated that acceptance and commitment therapy can have an effect on the anxiety levels of mothers with children with insecure attachment. This finding is consistent with previous studies by (Oskis et al., 2013) and (Pace et al., 2015). The explanation for the effect of acceptance and commitment therapy on anxiety aligns with the findings of other previous studies such as (Bahreini et al., 2024; Perry et al., 2017; Roshani et al., 2017; Babaei et al., 2015), and (Snow et al., 2005). It suggests that acceptance and commitment therapy helps individuals create a meaningful life while accepting the inevitable suffering and committing to live with it. When a person prepares themselves to build such a life, they may encounter various obstacles in the form of unwanted internal experiences, including thoughts, imaginations, feelings, bodily sensations, impulses, and memories. Acceptance and commitment therapy teaches effective mindfulness skills to manage these inner experiences. Mindfulness involves consciously and accepting directing our awareness to the present moment. These techniques help us live in the present, fully engage in our tasks instead of getting lost in thoughts, and allow our emotions to be as they are, coming and going, instead of trying to control them. By accepting our inner experiences, even painful memories, emotions, thoughts, and bodily sensations appear less threatening and unbearable. Thus, mindfulness assists in changing our relationship with these painful thoughts and feelings, reducing their impact and influence on our lives. These interventions focus on two fundamental processes: accepting unwanted inner experiences that are beyond our control and committing to live a valuable life and

taking action towards it.

The results have indicated that acceptance and commitment therapy can have a positive effect on the alexithymia of mothers with children exhibiting insecure attachment. This finding aligns with the results of previous studies, such as (Suslow & Junghanns, 2002), (Thorberg et al., 2009), and (Vanheule et al., 2007). The effect of acceptance and commitment therapy on the alexithymia of the sample individuals can be attributed to the adaptability of its concepts and interventions, which can be applied both individually and in group settings, as well as in counseling and psychotherapy centers. This finding is consistent with the results of other previous studies, such as (Vøllestad et al., 2011; Zeidner et al., 2012), and (Taylor & Bagby, 2004).

Acceptance and commitment therapy aims to reconcile the needs of the mother and the child, bridging the gap in their assessments. When the mother and child share a similar assessment of their life resources and demands, it strengthens family cohesion, forming the basis for improving family life quality and psychological flexibility. By focusing on enhancing intrapersonal skills (such as emotional and thought awareness) and interpersonal skills (communication skills and effective communication), acceptance and commitment therapy helps boost the psychological flexibility of the individuals in the sample.

The reduction of alexithymia through acceptance and commitment therapy is explained by the technique of acceptance or willingness to experience difficulty without attempting to suppress it, leading to a greater capacity to confront the challenges of personal, family, and social life. This approach diminishes avoidance, distress, fear of challenges, and ultimately, alexithymia. The primary objective of acceptance and commitment therapy is to cultivate and enhance flexibility. The ability to choose a more suitable option from among different choices enhances psychological well-being and a sense of relaxation, aiding individuals in coping adaptively with sources of stress. This treatment assists individuals in recognizing and expressing feelings, thereby reducing alexithymia. Acceptance and commitment therapy encourages the client to shift focus from symptom reduction to living a meaningful life.

Conclusion

The results indicated that the levels of anxiety and alexithymia underwent significant changes following the

intervention of ACT. ACT promoted a rich and meaningful life by accepting inevitable suffering and committing to living with it. When one prepared to build such a life, they encountered various obstacles in the form of undesirable internal experiences, including thoughts, imaginations, feelings, bodily sensations, impulses, and memories. ACT equipped individuals with effective mindfulness skills to manage these inner experiences. However, caution should be exercised when generalizing the findings of this study to other mothers, given that it was conducted solely on the population of mothers with children exhibiting an insecure attachment style.

The limitations of the present study were the quasi-experimental nature and the lack of easy access to mothers with children with insecure attachment. Counseling and psychotherapy centers should consider implementing these methods, particularly acceptance and commitment therapy, more widely and with greater focus. Future researchers should pay attention to the design and validation of scales that investigate the psychological and behavioral structures of families. Self-help booklets based on the exercises of this treatment method are useful for families. Cultural programs and educational workshops can increase families' awareness of acceptance and commitment, improve parenting, and promote the development of secure attachment styles.

Acknowledgments

The authors express their gratitude and appreciation to all participants.

Declaration of Interest

The authors of this article declared no conflict of interest.

Ethical Considerations

The study protocol adhered to the principles outlined in the Helsinki Declaration, which provides guidelines for ethical research involving human participants. Ethical considerations in this study were that participation was entirely optional.

Transparency of Data

In accordance with the principles of transparency and open research, we declare that all data and materials used in this study are available upon request.

Funding

This research was carried out independently with personal funding and without the financial support of any governmental or private institution or organization.

Authors' Contributions

All authors equally contribute to this study.

References

- Abdi Zarrin, S., & Nikkhah Siruei, Z. (2021). Psychological well-being: the role of self-differentiation, Alexithymia and ego-strength. *Shenakht Journal of Psychology and Psychiatry*, 8(3), 13-24. <https://doi.org/10.32598/shenakht.8.3.13>
- Akbari, M., & Elmî, R. (2017). Herpes simplex virus and human papillomavirus coinfections in hyperimmunoglobulin E syndrome presenting as a conjunctival mass lesion. *Case reports in medicine*, 2017(1), 1650841. <https://doi.org/10.1155/2017/1650841>
- Askari, M., & Karami, H. (2024). On the Relationship between Sensory Learning Styles and Reading Subskill Profiles: An Application of Fusion Model. *Language Related Research*, 15(3), 245-274. <https://doi.org/10.2139/ssrn.4873700>
- Babaei, S., Varandi, S. R., Hatami, Z., & Gharechahi, M. (2015). Metacognition beliefs and general health in predicting alexithymia in students. *Global journal of health science*, 8(2), 117. <https://doi.org/10.5539/gjhs.v8n2p117>
- Bagby, R. M., Taylor, G. J., & Parker, J. D. (1994). The twenty-item Toronto Alexithymia Scale—II. Convergent, discriminant, and concurrent validity. *Journal of psychosomatic research*, 38(1), 33-40. [https://doi.org/10.1016/0022-3999\(94\)90006-X](https://doi.org/10.1016/0022-3999(94)90006-X)
- Bahreini, F., Azizi, A., & Roohafza, H. (2024). Effectiveness of Bioenergy Economy-based Health Improvement versus Mindfulness-based Stress Reduction on the Occupational Stress and Psychosomatic Symptoms of Distressed Employees? *International Journal of Body, Mind & Culture* (2345-5802), 11(2). <https://doi.org/10.22122/ijbmc.v11i2.691>
- Beck, A. T., Epstein, N., Brown, G., & Steer, R. A. (1988). An inventory for measuring clinical anxiety: psychometric properties. *Journal of consulting and clinical psychology*, 56(6), 893. <https://doi.org/10.1037/0022-006X.56.6.893>
- Byrne, G., Ghráda, Á. N., O'Mahony, T., & Brennan, E. (2021). A systematic review of the use of acceptance and commitment therapy in supporting parents. *Psychology and Psychotherapy: Theory, Research and Practice*, 94, 378-407. <https://doi.org/10.1111/papt.12282>
- Çiçek Gümüş, E., & Öncel, S. (2023). Effects of acceptance and commitment therapy-based interventions on the mental states of parents with special needs children: Randomized controlled trial. *Current Psychology*, 42(23), 19429-19442. <https://doi.org/10.1007/s12144-022-03760-1>

- Darvishinia, N. (2023). AI in education: Cracking the code through challenges: A content analysis of one of the recent issues of Educational Technology and Society (ET&S) Journal. *Partners Universal International Innovation Journal*, 1(4), 61-71. <https://doi.org/10.5281/zenodo.826426>
- Eskandari, F., Aali, M., Hadisadegh, S. N., & Azadeh, M. (2024). Advances in breast cancer research using CRISPR/Cas9 system. *Nano Select*, 5(10), 2400015. <https://doi.org/10.1002/nano.202400015>
- Feyzabadi, Z. N., Nickzad, Z., Rejali, M., Behshahrian, A., & Amanat, A. (2025). Pessimism about Marriage as a Mediator Between Insecure Attachment and Marriage Age in Children of Divorced Parents: A Structural Equation Model. *International Journal of Body, Mind & Culture*, 12(6), 61-70. <https://doi.org/10.61838/ijbmc.v12i6.1007>
- Gould, E. R., Tarbox, J., & Coyne, L. (2018). Evaluating the effects of acceptance and commitment training on the overt behavior of parents of children with autism. *Journal of contextual behavioral science*, 7, 81-88. <https://doi.org/10.1016/j.jcbs.2017.06.003>
- Hahs, A. D., Dixon, M. R., & Paliliunas, D. (2019). Randomized controlled trial of a brief acceptance and commitment training for parents of individuals diagnosed with autism spectrum disorders. *Journal of contextual behavioral science*, 12, 154-159. <https://doi.org/10.1016/j.jcbs.2018.03.002>
- Hayes, S. C., Luoma, J. B., Bond, F. W., Masuda, A., & Lillis, J. (2006). Acceptance and commitment therapy: Model, processes and outcomes. *Behaviour research and therapy*, 44(1), 1-25. <https://doi.org/10.1016/j.brat.2005.06.006>
- Jiang, X., Sun, J., Song, R., Wang, Y., Li, J., & Shi, R. (2024). Acceptance and commitment therapy reduces psychological distress in patients with cancer: a systematic review and meta-analysis of randomized controlled trials. *Frontiers in Psychology*, 14, 1253266. <https://doi.org/10.3389/fpsyg.2023.1253266>
- Kappenberg, E. S., & Halpern, D. F. (2006). Kinship Center Attachment Questionnaire: Development of a caregiver-completed attachment measure for children younger than 6 years. *Educational and Psychological Measurement*, 66(5), 852-873. <https://doi.org/10.1177/0013164405285545>
- Karimian, Z., Haghayegh, S. A., Emami-Najafi-Dehkordi, S. M. H., & Raisi, M. (2023). The Comparison of the Effectiveness of Trans-diagnostic Treatment and the Acceptance and Commitment Therapy (ACT) on Quality of Life in Patients with Irritable Bowel Syndrome (IBS). *Journal of Sabzevar University of Medical Sciences*, 30(1), 22-34. <https://doi.org/10.30468/jsums.2023.1552>
- Loas, G., Baelde, O., & Verrier, A. (2015). Relationship between alexithymia and dependent personality disorder: a dimensional analysis. *Psychiatry research*, 225(3), 484-488. <https://doi.org/10.1016/j.psychres.2014.11.062>
- Lumley, M. A., Gustavson, B. J., Partridge, R. T., & Labouvie-Vief, G. (2005). Assessing alexithymia and related emotional ability constructs using multiple methods: interrelationships among measures. *Emotion*, 5(3), 329. <https://doi.org/10.1037/1528-3542.5.3.329>
- Malik, S., Wells, A., & Wittkowsky, A. (2015). Emotion regulation as a mediator in the relationship between attachment and depressive symptomatology: A systematic review. *Journal of affective disorders*, 172, 428-444. <https://doi.org/10.1016/j.jad.2014.10.007>
- Marino, F., Failla, C., Chilà, P., Minutoli, R., Puglisi, A., Arnao, A. A., Pignolo, L., Presti, G., Pergolizzi, F., & Moderato, P. (2021). The effect of acceptance and commitment therapy for improving psychological well-being in parents of individuals with autism spectrum disorders: a randomized controlled trial. *Brain sciences*, 11(7), 880. <https://doi.org/10.3390/brainsci11070880>
- Marsero, S., Ruggiero, G., Scarone, S., Bertelli, S., & Sassaroli, S. (2011). The relationship between alexithymia and maladaptive perfectionism in eating disorders: A mediation moderation analysis methodology. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 16(3), e182-e187. <https://doi.org/10.1007/BF03325130>
- MEHRABIZADEH, H. M., Taghavi, S., & ATARI, Y. A. (2009). Effect of group assertive training on social anxiety, social skills and academic performance of female students. https://www.behavsci.ir/article_67620.html
- Mikulincer, M., & Shaver, P. R. (2012). An attachment perspective on psychopathology. *World psychiatry*, 11(1), 11-15. <https://doi.org/10.1016/j.wpsyc.2012.01.003>
- Moghadam, H. I. (2024). Investigating the Relationship between Parenting Styles and Suicide Ideation with the Mediation of Avoidant Insecure Attachment Style. *International Journal of Body, Mind & Culture*, 11(2), 171-179.
- Monfaredi, Z., Malakouti, J., Farvareshi, M., & Mirghafourvand, M. (2022). Effect of acceptance and commitment therapy on mood, sleep quality and quality of life in menopausal women: a randomized controlled trial. *BMC psychiatry*, 22(1), 108. <https://doi.org/10.1186/s12888-022-03768-8>
- Oskis, A., Clow, A., Hucklebridge, F., Bifulco, A., Jacobs, C., & Loveday, C. (2013). Understanding alexithymia in female adolescents: The role of attachment style. *Personality and Individual Differences*, 54(1), 97-102. <https://doi.org/10.1016/j.paid.2012.08.023>
- Pace, C. S., Cavanna, D., Guiducci, V., & Bizzi, F. (2015). When parenting fails: alexithymia and attachment states of mind in mothers of female patients with eating disorders. *Frontiers in Psychology*, 6, 1145. <https://doi.org/10.3389/fpsyg.2015.01145>
- Perry, R. E., Blair, C., & Sullivan, R. M. (2017). Neurobiology of infant attachment: attachment despite adversity and parental programming of emotionality. *Current opinion in psychology*, 17, 1-6. <https://doi.org/10.1016/j.copsyc.2017.04.022>
- Roshani, F., Najafi, M., Naqshbandi, S., & Malekzade, P. (2017). Comparison of alexithymia in individuals with and without attention deficit/hyperactivity disorder. *Journal of Clinical Psychology*, 9(2), 73-82. <https://doi.org/10.22075/jcp.2017.11204.1109>
- Rostamifar, R., & Sajjadi, P. (2017). Investigation Role of Thought Control, Mindfulness, Distress Tolerance, in Prediction Signs of Borderline Personality Disorder in Female Students of Three Public High Schools in Golpayegan City. *International Journal of Health Sciences(II)*, 6989-7003. <https://doi.org/10.53730/ijhs.v6nS2.6680>
- Snow, M., Sullivan, K., Martin, E., & Helm, H. (2005). The adult scale of parental attachment: Psychometric properties, factor analysis, and multidimensional scaling. *Unpublished manuscript, Department of Leadership and Counselor Education, The University of Mississippi, University, Mississippi*.
- Suslow, T., & Junghanns, K. (2002). Impairments of emotion situation priming in alexithymia. *Personality and Individual Differences*, 32(3), 541-550. [https://doi.org/10.1016/S0191-8869\(01\)00056-3](https://doi.org/10.1016/S0191-8869(01)00056-3)
- Taylor, G. J., & Bagby, R. M. (2004). New trends in alexithymia research. *Psychotherapy and psychosomatics*, 73(2), 68-77. <https://doi.org/10.1159/000075537>
- Thorberg, F. A., Young, R. M., Sullivan, K. A., & Lyvers, M. (2009). Alexithymia and alcohol use disorders: A critical review. *Addictive behaviors*, 34(3), 237-245. <https://doi.org/10.1016/j.addbeh.2008.10.016>

- Vanheule, S., Desmet, M., Meganck, R., & Bogaerts, S. (2007). Alexithymia and interpersonal problems. *Journal of Clinical Psychology*, 63(1), 109-117. <https://doi.org/10.1002/jclp.20324>

Vøllestad, J., Sivertsen, B., & Nielsen, G. H. (2011). Mindfulness-based stress reduction for patients with anxiety disorders: Evaluation in a randomized controlled trial. *Behaviour research and therapy*, 49(4), 281-288. <https://doi.org/10.1016/j.brat.2011.01.007>

Wijk, I., Amsberg, S., Johansson, U.-B., Livheim, F., Toft, E., & Anderbro, T. (2023). Impact of an Acceptance and Commitment Therapy programme on HbA1c, self-management and psychosocial factors in adults with type 1 diabetes and elevated HbA1c levels: a randomised controlled trial. *BMJ open*, 13(12), e072061. <https://doi.org/10.1136/bmjopen-2023-072061>

Wiltgen, A., Arbona, C., Frankel, L., & Frueh, B. C. (2015). Interpersonal trauma, attachment insecurity and anxiety in an inpatient psychiatric population. *Journal of anxiety disorders*, 35, 82-87. <https://doi.org/10.1016/j.janxdis.2015.07.010>

Zeidner, M., Matthews, G., & Roberts, R. D. (2012). *What we know about emotional intelligence: How it affects learning, work, relationships, and our mental health*. MIT press. [https://books.google.com/books?id=7n7icwxvd2sC&lpg=PR7&ots=yx8ESXbcP8&dq=Zeidner%20M.%20Matt%20hews%20G.%20%26%20Roberts%20R.%20D.%20\(2012\).%20What%20we%20know%20about%20emotional%20intelligence%3A%20How%20it%20affects%20learning%20work%2C%20relationships%2C%20and%20our%20mental%20health.%20MIT%20press.%20&lr&pg=PR7#v=onepage&q&f=false](https://books.google.com/books?id=7n7icwxvd2sC&lpg=PR7&ots=yx8ESXbcP8&dq=Zeidner%20M.%20Matt%20hews%20G.%20%26%20Roberts%20R.%20D.%20(2012).%20What%20we%20know%20about%20emotional%20intelligence%3A%20How%20it%20affects%20learning%20work%2C%20relationships%2C%20and%20our%20mental%20health.%20MIT%20press.%20&lr&pg=PR7#v=onepage&q&f=false)

Zhao, B., Wang, Q., Wang, L., Chen, J., Yin, T., Zhang, J., Cheng, X., & Hou, R. (2023). Effect of acceptance and commitment therapy for depressive disorders: a meta-analysis. *Annals of general psychiatry*, 22(1), 34. <https://doi.org/10.1186/s12991-023-00462-1>