

Article type:
Original Research

1 Psychiatric Mental Health Nursing Department,
Faculty of Nursing, University of Baghdad, Iraq.

Corresponding author email address:
imanh@conursing.uobaghdad.edu.iq



Article history:

Received 21 May 2025
Revised 14 June 2025
Accepted 24 June 2025
Published online 30 July 2025

How to cite this article:

Alwan, I. H. (2025). Factors Influencing Stigma Toward Mental Illness Among Students at the University of Baghdad. *International Journal of Body, Mind and Culture*, 12(5), 112-121.



© 2025 the authors. This is an open-access article under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

Introduction

According to the World Health Organization (WHO), 450 million people worldwide suffer from mental illnesses, and one in every four people will be affected at some point in their lives. This stigma can have a negative influence on the lives of people with mental disorders, such as judgment, social isolation, and an increased risk of suicide. Stigma towards mental disorders can be defined as negative and discriminatory attitudes

Factors Influencing Stigma Toward Mental Illness Among Students at the University of Baghdad

Iman Hussein. Alwan¹ 

ABSTRACT

Objective: This study aimed to identify and analyze the key factors contributing to mental illness stigma among students at the University of Baghdad in Baghdad, Iraq.

Methods and Materials: A cross-sectional, descriptive study was conducted to achieve the study's objectives. Study Setting: The study was conducted among university students in Baghdad City, Iraq. A convenience sample of 300 students was included in the study. Study Tools: Part 1: Socio-demographic characteristics, including age, level of education, monthly family income, and number of siblings. Part 2: Quality of life was assessed using the World Health Organization Quality of Life (WHOQOL-BREF; World Health Organization, 2004). Part 3: The Stigma Scale was used to measure the feelings and experiences of individuals who have faced psychological distress, particularly focusing on societal reactions to their condition.

Findings: The mean age of the students was 22.21 ± 3.540 years, indicating that most participants were in their early twenties. Females (61.7%) constituted a larger proportion of the sample compared to males (38.3%). The mean score for mental illness stigma was 48.79 ± 6.811 , reflecting the overall stigma perceived by students. Among the influencing factors, the overall factors had the highest mean score (84.96 ± 14.439).

Conclusion: The study findings indicate that psychological well-being, overall health, and strong social support play a crucial role in reducing mental illness stigma among University of Baghdad students, with psychological factors having the most significant impact. These results highlight the importance of comprehensive strategies to promote mental health and effectively reduce stigma.

Keywords: Factors Influencing, Mental Illness Stigma, Nursing students.

towards people with mental disorders. This stigma can come from various sources, including family, friends, society, and mass media. The stigma against mental disorders in Indonesia has become a serious concern for the government and society (Davies, 2022).

Numerous universities around the world have implemented admissions policies to support students with disabilities and foster an inclusive learning environment. Sustainable Development Goal 3 (SDG3)

emphasizes the importance of mental health as a critical component of overall well-being at all stages of life, which is consistent with these policies (Davies, 2022).

While poor mental health creates difficulties in day-to-day living and impacts one's emotional, social, financial, and physical well-being, good mental health empowers people to make constructive contributions to society. Consequently, mental health continues to be a major global public health concern (Liu et al., 2024). Research has shown that younger college students are more likely than older students to suffer from mental health problems, which emphasizes the need for increased support in educational settings (Moudatsou et al., 2020; Staiger et al., 2018). In social psychiatry, the term stigma refers to a set of negative factors associated with individuals with a mental disorder. Specifically, according to the World Health Organization, stigma represents "a mark of shame, disgrace, or disapproval that leads to rejection, discrimination, and exclusion from social contexts and situations" (Ekman & Krasner, 2017; Thornicroft, 2018).

Attitudes regarding mental disease are greatly influenced by the stigma associated with mental health, especially among students. Numerous students have stigmatizing views about people with mental diseases, according to research (Ekman & Krasner, 2017). Such stigma can have a detrimental effect on the academic and social experiences of students who are dealing with mental health issues. One important determinant of the well-being of students with severe mental illness is peer interactions (Thornicroft, 2018). This emphasizes how student populations have a significant impact on how mental illnesses are treated and how society views them more broadly. Students who are sympathetic and inclusive are more likely to help those who suffer from mental illnesses and support initiatives that make it easier for them to get the care and services they need. Because of social, political, and cultural considerations, stigma can be a major deterrent for students who want to get help (Carrara et al., 2019).

Even though the number of students with disabilities has increased dramatically worldwide, they nevertheless face more obstacles in school than their peers without impairments. These difficulties restrict access to postsecondary education and career possibilities, and they also contribute to disparities in academic achievement (Lien et al., 2021). Eleven percent of

undergraduate students say they have a condition or impairment that limits their capacity to engage in social and academic activities fully. Students' quality of life (QOL) is significantly impacted by mental health disorders in particular (Holmqvist, 2020).

A person's physical health, social relationships, academic achievement, and general well-being are all impacted by psychotic disorders. Symptoms like delusions and hallucinations frequently cause social exclusion and misunderstanding. Patients' willingness to seek help is influenced by several factors, including stigma, attitudes, and the empathy of mental health experts, even in the face of excellent therapies (Gonzalez et al., 2017).

A higher level of internalized stigma was associated with greater perceived discrimination, lower social support, lower family income, older age, and distancing coping. These factors may be important to consider for developing tailored interventions in the future. (Yanos et al., 2001)

In patient care, empathy—the capacity to comprehend and experience another person's life on an emotional and cognitive level (Corrigan et al., 2014)—is essential. An empathic therapist can increase participation in treatment, promote hope, and lessen feelings of loneliness (Corrigan et al., 2014). According to research, patient outcomes improve and treatment effectiveness and quality of care are enhanced when mental health care workers have positive attitudes and higher levels of empathy (Dalky et al., 2020; Derksen et al., 2013). However, because people with psychotic disorders are frequently associated with negative stereotypes such as "unpredictable," "violent," or "aggressive," stigma among healthcare workers can still be problematic (Rössler, 2016). People may be deterred from obtaining therapy due to this stigma, as they fear discrimination (Koschorke et al., 2021).

Even among medical experts, psychotic disorders are one of the most stigmatized mental health problems, according to a systematic evaluation of 38 studies (Yanos et al., 2001). Although stigma is often lower among mental health professionals (Corrigan et al., 2014), this is frequently because they deal directly with patients (Derksen et al., 2013). On the other hand, professionals who experience associated stigma are more depersonalized, emotionally spent, and less satisfied with their jobs. Additionally, it may make patients feel

more ashamed of themselves, which would make their recuperation much more difficult (Dalky et al., 2020). Notwithstanding these results, it is still unknown how stigma and factors like age, occupation, education, and years of experience among mental health practitioners relate to one another (Yanos et al., 2001).

Cultural differences have a profound impact on people's perceptions and experiences with severe mental disorders (SMI), including schizophrenia. According to research, social and environmental factors influence both the risk and outcome of psychotic diseases. Individuals with schizophrenia in developing countries, such as Nigeria and India, frequently have better prognoses—characterized by lower mortality rates, fewer hospitalizations, and improved overall functioning—than those in developed countries, such as the United Kingdom and the United States (Rössler, 2016). Furthermore, non-affective acute remitting psychosis (NARP), a subtype of schizophrenia characterized by prolonged symptom-free intervals, is more prevalent in developing countries. This shows that cultural influences, specifically gene-environment interactions, have a significant impact on mental health outcomes. An aspect not to be underestimated is "diagnostic overshadowing," the underestimation of physical ailments in people with mental disorders (Jablensky & Sartorius, 2008). Without appropriate intervention measures, such as content moderation and psychological support, social media can become an echo chamber for these behaviors, exacerbating problems instead of providing a way out (Silva et al., 2017).

The presence of stigma among classmates frequently hinders students from getting the mental health care they require, owing to fear of being judged. Addressing the root causes of stigma and adopting interventions in student populations can help eliminate obstacles to mental health care (Lien et al., 2021). Fostering an open and tolerant campus environment can lead to improved support networks, allowing students to access needed mental health resources without fear of prejudice.

Universities can assist in creating a more inclusive environment that encourages students to seek aid, thereby increasing their academic and social well-being.

This study aims to assess the extent of mental disease stigma among students, providing insight into how they perceive mental health concerns and their prevalence.

2. Identify variables contributing to mental illness stigma among students. This entails identifying the key factors—whether psychological, social, or physical—that contribute to the stigma associated with mental illness.

3. Investigate the association between psychological, social, and physical aspects and the stigma of mental illness among students.

Methods and Materials

Study Design and Participants

A cross-sectional study, a descriptive design was used for this investigation. This design is great for determining the link between variables at a certain point in time, allowing for a full description of the amount of stigma and its associated components in the student population.

The study was carried out at the University of Baghdad in Iraq, which provides an important backdrop for understanding how university students in Iraq perceive and react to mental illness stigma.

The sample includes both male and female students, with a higher proportion of females (61.7%) compared to males (38.3%). Regarding marital status, the majority of participants are married (91.7%), while 7.9% are single and 0.3% are divorced. In terms of financial status, 70.6% report having sufficient income, while 29.4% find their income inadequate.

The majority of students (87.8%) are enrolled in morning study programs, whereas 12.2% attend evening studies. Regarding employment status, most students (69.0%) are full-time students, 25.7% are self-employed, and 5.3% are formally employed.

To determine the appropriate sample size, Slovin's formula was used:

$$n = N / (1 + Ne^2)$$

Where n represents the required sample size, N is the total student population, and e is the margin of error (set at 0.05). This method ensures that the sample is representative of the target population.

$$n = 1 + 1200 \times 0.05^2 / 1200 = 1 + 31200 = 41200 = 300$$

A non-probability convenience sampling technique was employed to pick 300 University of Baghdad students. This strategy provides a realistic and representative sample, but it may limit generalizability. The sample size is sufficient to yield relevant results.

Instruments

The Research and Ethics Committee approved the study involving students in the College of Nursing at the University of Baghdad. Tools include the interviewing questionnaire (Tool I).

The interviewing questionnaire was specifically created to collect data relevant to the study's aims. The questionnaire has two parts:

1. Part 1 includes socio-demographic information such as age, education level, monthly household income, number of siblings, and other important personal factors.

2. In Part 2, the WHOQOL-BREF scale (World Health Organization, 2004) was used to assess quality of life, which includes physical and psychological well-being.

Social relationships. Environmental factors.

- Respondents rated satisfaction on a 5-point Likert scale (1 = dissatisfied, 5 = fully satisfied). The scale's good internal consistency ($\alpha = .91$) indicates accurate measurement of quality of life.

- The Stigma Scale captures the feelings and experiences of persons suffering from psychiatric conditions, with a focus on society's reactions. This measure was developed through qualitative research, which included in-depth interviews with persons who had faced mental health issues. It comprises 28 questions, each of which is graded on a scale of 0 to 4, with higher scores indicating greater stigma.

- Questions are scored as "A" for agreement and "D" for disagreement.

- The scale evaluates the social and psychological effects of stigma, offering insights into societal views towards individuals with mental health disorders.

Participants are advised to respond based on their first perceptions, making sure to be honest in representing their experiences with stigma. This measure can assist in identifying areas for intervention and guide future efforts to reduce mental health stigma in society and healthcare settings.

A committee of five mental health nursing professionals from the College of Nursing at the University of Baghdad assessed the study tools to ensure

their validity. The tools were revised based on expert comments to ensure their relevance and correctness.

Cronbach's Alpha test was used to assess the study tools' internal consistency and dependability, and the reliability coefficient was 0.84. This suggests that the instruments are highly consistent and reliable. Cronbach's Alpha scores greater than 0.7 are deemed acceptable, and a coefficient of 0.84 indicates that the questionnaire accurately measures the target constructs. This level of reliability improves confidence in the accuracy of the collected data and supports the study's findings.

Data Analysis

During the preliminary phase, the researcher conducted a thorough evaluation of relevant literature, which included books, scholarly articles, and both local and international journals. The goal was to build upon existing knowledge and identify gaps in the literature.

This review helps to modify the study methodology and ensure that the research is anchored in existing knowledge of mental illness stigma.

Pilot Study.

A pilot study was conducted using 10% of the target sample (30 students) to determine the applicability and effectiveness of the research tools. The feedback from this pilot study was used to make required changes to the study instruments, ensuring that they are relevant and accurate for the full-scale study.

The fieldwork was carried out between October 2023 and January 2024. During this time, the researcher introduced themselves to the students of the University of Baghdad. The interview process included discussing each item on the questionnaire to verify that the students understood the questions and could reply correctly. Assistance was offered when needed, and each interview lasted around 30-40 minutes.

Findings and Results

The findings of the data analysis are presented systematically in the following (Table 1):

Table 1

Distribution of Students in the University of Baghdad by their Socio-demographic Variables (SDVs)

| SDVs | Classification | No. | % |
|-------------------|------------------------|-------------------|------|
| Age (years) | M \pm Std. Deviation | 22.21 \pm 3.540 | |
| Sex | Male | 116 | 38.3 |
| | Female | 187 | 61.7 |
| Marital Status | Single | 24 | 7.9 |
| | Married | 278 | 91.7 |
| | Divorced | 1 | 0.3 |
| Monthly Income | Enough | 214 | 70.6 |
| | Not Enough | 89 | 29.4 |
| Type of Study | Morning | 266 | 87.8 |
| | Evening | 37 | 12.2 |
| Employment Status | Employed | 16 | 5.3 |
| | Self-employed | 78 | 25.7 |
| | Students | 209 | 69.0 |

No. Number; %= Percentage; M= Mean; Std. Deviation = Standard deviation

Table 2 presents the distribution of students at the University of Baghdad based on their socio-demographic variables (SDVs). The mean age of the students is 22.21 ± 3.540 years, indicating that most participants are in their early twenties.

In terms of sex distribution, females (61.7%) constitute a larger proportion of the sample compared to males (38.3%). Regarding marital status, the majority of students are married (91.7%), while a small percentage are single (7.9%) or divorced (0.3%). In terms of

monthly income, 70.6% of students report having enough income, whereas 29.4% find their income insufficient.

The type of study is predominantly morning programs (87.8%), with a smaller portion enrolled in evening studies (12.2%). In terms of employment status, most students (69.0%) are full-time students, while 25.7% are self-employed, and a small fraction (5.3%) are formally employed.

Table 2

Distribution of Mental Illness Stigma and Its Influencing Factors among Students in the University of Baghdad

| Factors | Mean \pm Std. Deviation |
|-----------------------|---------------------------|
| Mental Illness Stigma | 48.79 \pm 6.811 |
| Physical Factors | 23.17 \pm 4.398 |
| Psychological Factors | 20.00 \pm 4.459 |
| Social Factors | 10.04 \pm 2.740 |
| Environmental Factors | 26.99 \pm 5.897 |
| Overall Factors | 84.96 \pm 14.439 |

M: Mean for total score, Std. Deviation =Standard Deviation for total score

Table 2 presents the distribution of mental illness stigma and its influencing factors among students at the University of Baghdad. The mean score for mental illness

stigma is 48.79 ± 6.811 , indicating a moderate level of stigma among students. Among the influencing factors, environmental factors have the highest mean ($26.99 \pm$

5.897), suggesting that environmental conditions play a significant role in shaping stigma. Physical factors have a mean of 23.17 ± 4.398 , while psychological factors have a slightly lower mean of 20.00 ± 4.459 , indicating their substantial influence. Social factors show the lowest

mean (10.04 ± 2.740), suggesting a relatively smaller but still notable contribution to stigma. The overall influencing factors have a mean score of 84.96 ± 14.439 , reflecting the combined impact of these variables on mental illness stigma among university students.

Table 3

Factors Influencing Mental Illness Stigma among Students in the University of Baghdad

| Variables | Unstandardized Coefficients | | Standardized Coefficients | T | Sig. |
|-----------------------|-----------------------------|------------|---------------------------|--------|------|
| | B | Std. Error | Beta | | |
| Physical factors | -.305 | .088 | -.197 | -3.487 | .001 |
| Psychological factors | -.457 | .084 | -.299 | -5.438 | .000 |
| Social factors | -.223 | .052 | -.241 | -4.316 | .000 |
| Environmental factors | -.049 | .065 | -.016 | -0.836 | .079 |
| Overall Factors | -.135 | .026 | -.287 | -5.189 | .000 |


Dependent Variable: Mental Illness Stigma

Table 3 presents the factors influencing mental illness stigma among students at the University of Baghdad. Among these factors, psychological aspects have the strongest negative association with mental illness stigma ($B = -0.457$, $Beta = -0.299$, $T = -5.438$, $p < 0.001$), indicating that better psychological well-being significantly reduces stigma. Overall factors also show a substantial negative influence ($B = -0.135$, $Beta = -0.287$, $T = -5.189$, $p < 0.001$), suggesting that a combination of various determinants contributes to lowering stigma levels. Additionally, social aspects demonstrate a significant negative relationship ($B = -0.223$, $Beta = -0.241$, $T = -4.316$, $p < 0.001$), implying that stronger social support plays a key role in reducing stigma. Lastly, physical aspects have the least impact but remain significant ($B = -0.305$, $Beta = -0.197$, $T = -3.487$, $p = 0.001$), highlighting the role of physical health in influencing stigma.

The figures illustrate the negative relationships between various influencing factors and mental illness stigma among university students. Figure 1 depicts the association between physical factors and stigma, showing a moderate negative effect ($R^2 = 0.0388$), suggesting that improved physical well-being contributes to reduced stigma. Figure 2 presents the strongest negative correlation, highlighting the impact of psychological factors ($R^2 = 0.0895$), indicating that better psychological health significantly lowers stigma. Figure 3 demonstrates the role of social factors ($R^2 = 0.0583$), emphasizing that stronger social support is linked to lower stigma levels. Finally, Figure 4 represents the overall influence of all factors combined ($R^2 = 0.0821$), reinforcing the idea that a holistic approach incorporating physical, psychological, and social well-being plays a crucial role in reducing mental illness stigma.

Figure 1

A) Physical Factors & Mental Illness Stigma B) Psychological Factors & Mental Illness Stigma C) Social Factors & Mental Illness Stigma D) Overall Factors & Mental Illness Stigma



Discussion and Conclusion

Our study's socio-demographic characteristics share numerous similarities with those investigated in "Perceived Stigma Towards Mental Illness Among College Students of Western Nepal" by Pokharel and Pokharel (2017). Both studies acknowledge the impact of gender, marital status, wealth, and employment on perceptions of mental illness stigma. In all research, gender and socioeconomic level were identified as significant factors in stigma, while marital status and employment also played crucial roles in determining views toward mental health. These findings indicate that the impact of socio-demographic characteristics on

stigma may be similar across contexts, emphasizing the importance of culturally responsive approaches to addressing mental health stigma.

In addition to socio-demographic factors, Pokharel & Pokharel (2017) underline the role of awareness and education in reducing mental health stigma. They discovered that those with higher levels of education have more positive attitudes regarding mental health, as opposed to people who know little about mental health disorders (Pokharel & Pokharel, 2017). This finding supports the value of educational interventions in reducing stigma, since increasing information can break down preconceptions and build empathy for persons with mental health issues.

. Furthermore, Pokharel & Pokharel emphasized that personal or family experience with mental illness is important in developing perspectives. According to their findings, pupils who had personal or familial experiences with mental health concerns were more likely to have a favorable attitude regarding mental illness. This finding emphasizes the need to encourage open discussions about mental health, as personal exposure to mental illness appears to play a major role in eliminating stigma. These findings are especially important to our study because they highlight the importance of including educational programs and fostering conversation about mental health to confront and change deeply ingrained stigmatizing attitudes.

Stigma against mental disorders is a critical issue, given, on the one hand, its influence on the well-being and quality of life of people with mental illnesses (Dalky et al., 2020; Jablensky et al., 1992; Koschorke et al., 2021; Linney et al., 2020; Rössler, 2016; Valery & Prouteau, 2020), and on the other. To our knowledge, there is currently no multi-component program aimed at reducing stigma associated with mental diseases. The findings of this study are consistent with those of (Huang, 2021), demonstrating the strong impact of mental illness stigma on individuals, albeit with differences in the elements that have

In contrast, the previous study identified emotional and behavioral factors as the most strongly correlated with reduced quality of life among patients. It revealed moderate to large negative correlations between the Affect and Behavior domains of the SSS-S scale and all domains of the WHOQOL-BREF ($r = -0.634$ to -0.741 , $p < 0.01$). The study found that cognitive elements had no significant impact on quality of life. Physical and psychological components had lower mean scores (23.17 ± 4.398 and 20.00 ± 4.459 , respectively) than social factors.

These data indicate that social factors are more important in generating stigma among students. In contrast, emotional and behavioral elements appear to have a greater impact on quality of life among patients. This disparity may reflect the differences in life experiences between students and those with persistent mental disorders, with students being more impacted by social beliefs and patients being more affected by their emotional and behavioral experiences related to the condition.

As a result, this research lends support to the idea that mental illness stigma spreads beyond patients to other populations, such as students. However, the factors driving stigma differ, emphasizing the importance of focused interventions that address the most significant concerns for each group. Improving social support may be important for students, whilst addressing emotional and behavioral components is critical for patients' quality of life.

The study "Stigma of Psychological Disorder among University Students" sheds light on the elements that influence mental illness stigma among students at the University of Baghdad. The study found a substantial negative correlation ($B = -0.457$, $Beta = -0.299$, $T = -5.438$, $p < 0.001$) between psychological well-being and mental illness stigma. Social elements have a substantial negative connection ($B = -0.223$, $Beta = -0.241$, $T = -4.316$, $p < 0.001$), indicating the need for strong social support in minimizing stigma. Physical elements had a substantial impact ($B = -0.305$, $Beta = -0.197$, $T = -3.487$, $p = 0.001$), emphasizing the importance of physical health in determining stigma.

Our study's socio-demographic characteristics share numerous similarities with those investigated in "Perceived Stigma Towards Mental Illness Among College Students of Western Nepal" by Pokharel and Pokharel (2017). Both studies acknowledge the impact of gender, marital status, wealth, and employment on perceptions of mental illness stigma (Pokharel & Pokharel, 2017). In all research, gender and socioeconomic level were identified as significant factors in stigma, while marital status and employment also played crucial roles in determining views toward mental health. These findings indicate that the impact of socio-demographic characteristics on stigma may be similar across contexts, emphasizing the importance of culturally responsive approaches to addressing mental health stigma.

These studies demonstrate the multiple nature of mental illness stigma and the need to address psychological, social, and physical issues to minimize stigma and improve quality of life. Interventions aimed at these areas may be useful in reducing stigma and enhancing well-being among people with mental health issues.

The stigma associated with mental illness remains a significant barrier to student well-being and access to

mental health treatments. This study examines how psychological, social, and cultural factors influence stigma, resulting in presumptions and a reluctance to seek treatment. To address stigma, educational initiatives must foster understanding, empathy, and inclusivity. Universities play an important role in creating a supportive atmosphere for students, thereby increasing their mental health and societal acceptance of mental illness.

Acknowledgments

We want to express our gratitude to the dedicated students of Al University of Baghdad who participated in this study.

Declaration of Interest

The authors of this article declared no conflict of interest.

Ethical Considerations

The study protocol adhered to the principles outlined in the Helsinki Declaration, which provides guidelines for ethical research involving human participants. Ethical considerations in this study were that participation was entirely optional.

Transparency of Data

By the principles of transparency and open research, we declare that all data and materials used in this study are available upon request.

Funding

This research was carried out independently with personal funding and without the financial support of any governmental or private institution or organization.

Authors' Contributions

All authors equally contribute to this study.

References

- Carrara, B. S., Ventura, C. A. A., Bobbili, S. J., Jacobina, O. M. P., Khenti, A., & Mendes, I. A. C. (2019). Stigma in health professionals towards people with mental illness: An integrative review. *Arch Psychiatr Nurs*, 33(4), 311-318. <https://doi.org/10.1016/j.apnu.2019.01.006>
- Corrigan, P. W., Druss, B. G., & Perlick, D. (2014). The impact of mental illness stigma on seeking and participating in mental health care. *Psychological Science in the Public Interest*, 15(2), 37-37. <https://doi.org/10.1177/1529100614531398>
- Dalky, H. F., Abu-Hassan, H. H., Dalky, A. F., & Al-Delaimy, W. (2020). Assessment of mental health stigma components of mental health knowledge, attitudes, and behaviors among Jordanian healthcare providers. *Community Ment Health J*, 56(3), 524-531. <https://doi.org/10.1007/s10597-019-00509-2>
- Davies, J. (2022). Analysis of Factors Influencing Stigma Against Mental Disorders in Indonesia: Systematic Literature Review. *Scientific Journal of Psychology*, 5(1), 165. <https://doi.org/10.37275/scipsy.v5i1.165>
- Derksen, F., Bensing, J., & Lagro-Janssen, A. (2013). Effectiveness of empathy in general practice: a systematic review. *Br J Gen Pract*, 63(606), e76-e84. <https://doi.org/10.3399/bjgp13X660814>
- Ekman, E., & Krasner, M. (2017). Empathy in medicine: Neuroscience, education, and challenges. *Medical Teacher*, 39(2), 164-173. <https://www.tandfonline.com/doi/abs/10.1080/0142159x.2016.1248925>
- Gonzalez, H. M., Tarraf, W., Whitfield, K. E., & Vega, W. A. (2017). The Epidemiology of Mental Disorders in Adults. *The American Journal of Geriatric Psychiatry*, 25(4), 255-266. <https://pubmed.ncbi.nlm.nih.gov/9285988/>
- Holmqvist, R. (2020). Associations between staff feelings toward patients and treatment outcome at psychiatric treatment homes. *J Nerv Ment Dis*, 188(6), 366EP - 371. <https://doi.org/10.1097/00005053-200006000-00007>
- Huang, W. Y. (2021). Stigma of psychological disorders and quality of life among individuals with mental illness: A community-based program. *Journal name*, 0(0), 0-0. <https://pmc.ncbi.nlm.nih.gov/articles/PMC11262073/>
- Jablensky, A., & Sartorius, N. (2008). The epidemiology of schizophrenia. *The Schizophrenia Bulletin*, 34(3), 409-414. <https://journals.sagepub.com/doi/abs/10.1177/070674379704200204>
- Jablensky, A., Sartorius, N., Korten, A. E., & Phillips, L. J. (1992). Psychotic disorders in urban areas: an overview of the WHO Collaborative Study on Psychoses. *Psychological medicine*, 22(3), 647-657. <https://journals.sagepub.com/doi/abs/10.1080/j.1440-1614.2000.00728.x>
- Koschorke, M., Oexle, N., Ouali, U., Cherian, A. V., Deepika, V., & Mendon, G. B. (2021). Perspectives of healthcare providers, service users, and family members about mental illness stigma in primary care settings: A multi-site qualitative study of seven countries in Africa, Asia, and Europe. *PLoS One*, 16(10), e0258729. <https://doi.org/10.1371/journal.pone.0258729>
- Lien, Y., Lin, H., Lien, Y., Tsai, C., Wu, T., & Li, H. (2021). Challenging mental illness stigma in healthcare professionals and students: a systematic review and network meta-analysis. *Psychol Health*, 36(6), 669-684. <https://doi.org/10.1080/08870446.2020.1828413>
- Linney, C., Ye, S., Redwood, S., Mohamed, A., Farah, A., & Biddle, L. (2020). "Crazy person is crazy person. It doesn't differentiate": an exploration into Somali views of mental health and access to healthcare in an established UK Somali community. *Int J Equity Health*, 19(1), 190. <https://doi.org/10.1186/s12939-020-01295-0>
- Liu, X., Yin, M., Li, Z., & Wang, D. (2024). Psychosocial Correlates of Internalized Stigma Among Chinese Individuals With Severe Mental Illness. *Journal of Psychosocial Nursing*

- and *Mental Health Services*, 62(1), 37-44.
<https://doi.org/10.3928/02793695-20230726-05>
- Moudatsou, M., Stavropoulou, A., Philalithis, A., Koukouli, S., Ekman, E., & Krasner, M. (2020). The role of empathy in health and social care professionals. Empathy in medicine: Neuroscience, education, and challenges. *Healthcare (Basel)*, 8(1), 26-173. <https://doi.org/10.3390/healthcare8010026>
 10.1080/0142159X.2016.1248925
- Pokharel, S., & Pokharel, R. (2017). Perceived stigma towards mental illness among college students of Western Nepal. *Journal name*, 2(3SP - 0), 0.
<https://doi.org/10.3126/bjhs.v2i3.18946>
- Rössler, W. (2016). The stigma of mental disorders: A millennia-long history of social exclusion and prejudices. *EMBO Rep*, 17(9), 1250-1253. <https://doi.org/10.15252/embr.201643041>
- Silva, R. D. D. C., Albuquerque, S. G. C., Muniz, A. D. V., Filho, P. P. R., Ribeiro, S., & Pinheiro, P. R. (2017). Reducing the schizophrenia stigma: a new approach based on augmented reality. *Comput Intell Neurosci*, 2017, 2721846-2721810.
<https://doi.org/10.1155/2017/2721846>
- Staiger, T., Waldmann, T., Oexle, N., Wigand, M., & Rüsch, N. (2018). Intersections of discrimination due to unemployment and mental health problems: the role of double stigma for job- and help-seeking behaviors. *Soc Psychiatry Psychiatr Epidemiol*, 53(10), 1091-1098.
<https://doi.org/10.1007/s00127-018-1535-9>
- Thornicroft, G. P. Y. (2018). Stigma and discrimination limit access to mental health care. *Epidemiol Psychiatr Soc*, 17(1), 14-19. <https://doi.org/10.1017/S1121189X00002621>
- Valery, K.-M., & Prouteau, A. (2020). Schizophrenia stigma in mental health professionals and associated factors: A systematic review. *Psychiatry research*, 290, 113068.
<https://www.sciencedirect.com/science/article/pii/S0165178120303024>
- Yanos, P. T., Primavera, L. H., & Rosenfeld, B. (2001). The role of social support in quality of life among individuals with schizophrenia. *Psychiatric Services*, 52(4), 455-459.
<https://doi.org/10.1176/appi.ps.52.4.493>